

Veterinary Package Insert

TP-ERYSIN 300MG/G POWDER (30% Erythromycin Thiocyanate)

PRODUCT DESCRIPTION

White to off-white colour powder, which contains 300mg of erythromycin thiocyanate (277.6mg of erythromycin) per gram in 1kg sachet.

PHARMACODYNAMICS & PHARMACOKINETICS

Erythromycin, an antibiotic of the macrolide group, is active against Gram-positive organisms (*Streptococcus spp.*, *Staphylococcus spp.*, *Corynebacterium spp.*, *Erysipelothrix rhusiopathiae*, *Listeria monocytogenes*, *Bacillus anthracis*, *Clostridium spp.*, *Actinomyces spp.*). Certain germs Gram negative (*Pasteurella multocida*, *Actinobacillus pleuropneumoniae*, *Haemophilus hemolyticus*) and mycoplasma. It is also active against leptospires.

The intestinal absorption of erythromycin after oral administration is good. Erythromycin is rapidly distributed in tissues; its weak base character allows wide tissue and intracellular diffusion, especially at the level of pulmonary alveoli. Erythromycin is actively excreted in the bile at significant amounts, which can be reabsorbed in the intestine. Only 5% of erythromycin is excreted in the urine.

INDICATION

In poultry:

- Treatment of infections due to germs sensitive to erythromycin.

TARGETED SPECIES

Poultry.

RECOMMENDED DOSAGE AND ADMINISTRATION

For oral administration.

20 mg erythromycin (72mg of product) per kg body weight per 24 hours in liquid food, milk or drinking water, for 3 days

Poultry:

Administer in drinking water at a rate of 0.72 g of powder per liter for 3 consecutive days on the basis of a water consumption of 100 ml per kg.

The amount of medicinal drinking water consumed by poultry depends on their physiological and clinical condition. In order to obtain the recommended dosage, the concentration of

erythromycin in the drinking water should be adjusted accordingly.

CONTRAINDICATIONS

Do not use in case of allergy to erythromycin or other macrolide.

WARNINGS & PRECAUTIONS.

Special precautions for use in animals:

Inappropriate use of the product may increase the prevalence of erythromycin-resistant bacteria.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

People with known hypersensitivity to macrolides should avoid contact with the veterinary medicinal product.

In case of reaction after exposure (rash for example), seek medical advice.

INTERACTIONS WITH OTHER MEDICATIONS

Not known.

PREGNANCY AND LACTATION

Studies in laboratory animals have not shown any teratogenic effects.

The safety of the specialty has not been studied in breeding poultry.

The specialty can only be used after evaluation of the benefit / risk ratio by the veterinarian.

ADVERSE EFFECTS

Not known.

OVERDOSE & TREATMENT

Not known.

WITHDRAWAL PERIOD

Poultry:

Meat and offal: 1 day.

Eggs: 0 days.

MAXIMUM RESIDUAL LIMIT (MRL)

Substance	Food	Maximum Residue Limits (MRLs) in µg/kg
Erythromycin	Edible offal, muscle, egg (mammalian and poultry)	300

STORAGE CONDITION

Store below 30°C. Protect from direct sunlight.

SHELF LIFE

2 years.

After 1st opening, reconstitution or dilution, use within 24 hours.

PACKING

100g

500g

1kg

THYE PHARMA SDN. BHD.

(manufacturer & product registration holder)

No.2 Lorong Industri Ringan Permatang

Tinggi 8,

Kawasan Industri Ringan Permatang Tinggi,

14000 Bukit Mertajam,

Pulau Pinang.

Revised October 2019