

Package Insert

Brand / product name	OC 200 g/kg Premix Powder
Name and Strength of active substance (s)	Composition in 1 kg: Oxytetracycline (as Dihydrate).....200 g (Equivalent 215.7 g of oxytetracycline dihydrate)
Product Description	Beige green powder
Pharmacodynamics	Oxytetracycline reversibly binds to the 30S ribosomal fraction receptors, leading to blocking of the aminoacyl-tRNA binding at the corresponding site of the ribosome-messenger RNA complex. This results in an inhibition of the protein synthesis and thus a stop of the growth of the bacterial culture. Oxytetracycline has a mainly bacteriostatic activity. The bacteriostatic activity of oxytetracycline involves penetration of the substance into the bacterial cell. The penetration of oxytetracycline is exerted by both passive and active diffusion. The main mode of possible resistance is related to the possible presence of an R factor responsible for a decrease in the active transport of oxytetracycline. Oxytetracycline is a broad-spectrum antibiotic. It is primarily active against gram-positive and negative microorganisms, aerobic and anaerobic, as well as against mycoplasma, chlamydia and Rickettsiae. Acquired resistance to oxytetracycline has been reported. Such resistance is usually of plasmid origin. Cross resistance to other tetracyclines is possible. Continuous treatment with low doses of oxytetracycline may also result in increased resistance to other antibiotics.
Pharmacokinetics	For most species, oxytetracycline is rapidly (2-4 h) absorbed after oral administration in fasting animals and has a bioavailability of 60% to 80%. This bioavailability can be decreased in the presence of food in the stomach because oxytetracycline forms insoluble chelates with the divalent or trivalent cations (Mg, Fe, Al, Ca) that they contain. In pigs, the influence of food is negligible on the bioavailability of oxytetracycline which is less than 5%. Oxytetracycline binds to plasma proteins in a variable manner depending on the species (20-40%). Its distribution is wide. Oxytetracycline diffuses throughout the body, with the highest concentrations found in the kidneys, liver, spleen and lungs. Oxytetracycline crosses the placental barrier. Oxytetracycline is eliminated unchanged, mainly by the urinary tract. It is also excreted by the bile duct but a large proportion of oxytetracycline is reabsorbed by the small intestine (enterohepatic cycle).
Environmental Properties	N/A

Package Insert

Indication	In lambs, kids (young goat), pigs, chickens, turkeys: - Treatment and prevention in an infected environment of respiratory infections and infections digestives due to germs sensitive to oxytetracycline.
Recommended Dosage	In pigs: Prevention in an infected environment: 25 mg of oxytetracycline per kg bodyweight and per day, orally, for 10 days. Treatment: 50 mg oxytetracycline per kg bodyweight and per day, orally, for 10 days. In lambs, kids (young goat), poultry: Prevention in infected medium: 20 mg of oxytetracycline per kg body weight per day, orally, for 10 days. Treatment: 40 mg of oxytetracycline per kg bodyweight and per day, orally, for 10 days. Incorporation rate: The amount of medicated feed consumed by the animals depends on their physiological (age) and clinical status. In order to respect the recommended doses of oxytetracycline and take into account the actual dietary intake, the rate of incorporation may be increased or possibly decreased to the minimum incorporation limit of 1 kg / tonne. This can lead to different ppm oxytetracycline levels in the medicated feed of those mentioned below for information purposes. Poultry: For a daily food intake of 100 g of feed per kg of live weight, this dosage corresponds, in prevention, to 200 ppm of oxytetracycline in the feed, ie an incorporation rate of 1 kg of medicated premix. per tonne of food, and in treatment, 400 ppm prevention of oxytetracycline in the food, a rate of incorporation of 2 kg of medicated premix per tonne of food. Pigs: For a daily dietary intake of 50 g of feed per kg of live weight, this dosage corresponds, in prevention, to 500 ppm of oxytetracycline in the feed, ie a incorporation rate of 2.5 kg of medicated premix. per ton of feed and 1000 ppm oxytetracycline in the feed, which is an incorporation rate of 5 kg of medicated premix per tonne of feed. Lambs and kids (young goat): For a daily food intake of 20 g of feed / kg of live weight, this dosage corresponds, in prevention, to 1000 ppm of oxytetracycline in the feed, ie a rate of incorporation of 5 kg of medicated premix. per tonne of feed and in treatment, prevention of oxytetracycline at 2000 ppm in the feed, ie a rate of incorporation of 10 kg of medicated premix per ton of feed.

Package Insert

Mode of Administration	Oral
Contraindication	Do not use in case of known allergy to oxytetracycline or any other substance of the tetracycline group. Do not use in case of known resistance to tetracyclines.
Warning and Precautions	a) Special warnings for each target species None. b) Special precautions for use i) Special precautions for use in animals This premix is intended for the manufacture of medicated feeds and can not be used as is; the rate of incorporation of the premix into the feed can not be less than 1 kg / tonne. (ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals Do not handle this product if you are allergic to tetracyclines. In case of reaction after exposure to the product (rash for example), consult a doctor. iii) Other precautions None.
Interaction with other Medicaments	Divalent or trivalent cations (Mg, Fe, Al, Ca) can chelate tetracyclines. Tetracyclines should not be administered with antacids, aluminum-based gels, mineral-based preparations as insoluble complexes are formed, which decreases the absorption of the antibiotic.
Treatment on usage during pregnancy and lactation	Oxytetracycline showed no evidence of embryotoxicity or teratogenicity in laboratory animals. In mammals, oxytetracycline passes the placental barrier, causing staining of teeth and slowing of fetal growth. Tetracyclines are found in breast milk. Product safety has not been evaluated in pregnant or lactating animals. The use of the product in pregnant or lactating animals should be subject to a risk / benefit assessment by the veterinarian.
MRL	<u>Fat:-</u> Sheep, pigs, chicken, turkeys – 10 µg/kg <u>Muscle:-</u> Sheep, pigs, chicken, turkeys – 100 µg/kg <u>Liver:-</u> Sheep, pigs, chicken, turkeys – 300 µg/kg

Package Insert

	<u>Kidney:-</u> Sheep, pigs, chicken, turkeys – 600 µg/kg <u>Eggs:-</u> 200 µg/kg
Adverse Effects/ undesirable Effects	As with all tetracyclines, adverse effects have been noted such as gastrointestinal disturbances and less frequently allergic reactions and photosensitivity.
Overdose and treatment	See section “Undesirable effects”.
Incompatibilities (for injection only)	N/A
Withdrawal period(s)	Meat and offal: 14 days. Eggs: zero days.
Storage conditions	Store below 30 °C (room temperature).
Dosage form and packaging available	Dosage form: Powder Packaging: 25 kg
Name and address of manufacturer	Weissen Company (M) Sdn. Bhd. 15 & 17, Jalan Metro Perdana Barat 12, Kepong Sri Edaran Ind. Park, 52100 Kuala Lumpur, Malaysia.
Name and address of registration holder	Weissen Company (M) Sdn. Bhd. Lot 105, No. 84, Jalan Velox 23, Taman Velox, 48000 Rawang, Selangor, Malaysia.
Date of revision of Package insert	02.10.2024
Shelf life of veterinary products as packaged for sales	24 months
Shelf life after reconstitution in feed	24 hours