

Side Effects:

Occasional side-effects such as nausea, abdominal discomfort, vomiting and diarrhoea may be experienced. Reversible hearing loss associated with doses of erythromycin usually greater than 4 g per day has been reported. Allergic reactions are rare and mild, although anaphylaxis has occurred extremely rarely. There are no reports implicating erythromycin products with abnormal tooth development and only rare reports of damage to the blood, kidneys, liver or central nervous system.

Symptoms and Treatment of Overdose:

Symptoms - Hearing loss, severe nausea, vomiting and diarrhoea.

Treatment - Gastric lavage, general supportive measures.

Storage Condition:

Keep container tightly closed. Store in a dry place (below 30°C). Protect from light. Once reconstituted the mixture should be kept in the refrigerator and used within one week of preparation.

Pack Size:**Ettrocin 250 Film-Coated Tablet**

Plastic container of 30's.

Blister pack of 1x10's, 10x10's, 50x10's and 100x10's.

Ettrocin for Suspension 200mg/5ml

Plastic bottles containing powder for the preparation of 60ml and 120ml suspension.

Products Registration Number:

Ettrocin 250 Film-Coated Tablet - MAL19860690AZ

Ettrocin for Suspension 200mg/5ml - MAL19860670AZ

ROYCE®**Product Holder / Manufactured by:**

ROYCE PHARMA MFG. SDN.BHD. (650435-X)

PT 1663, Nilai Industrial Estate,
71800 Nilai, Negeri Sembilan, Malaysia.

Revision date: 210218

ROYCE®

Ettrocin 250 Film-Coated Tablet
Ettrocin for Suspension 200mg/5ml
(Erythromycin Stearate / Erythromycin
Ethylsuccinate - Antibiotic)

Presentation:**Ettrocin 250 Film-Coated Tablet**

A round and red coloured film-coated tablet which is plain on both sides.

Ettrocin for Suspension 200mg/5ml

Bottle containing white powder for the preparation of 60ml. Orange flavoured, white suspension.

Content:**Ettrocin 250 Film-Coated Tablet**

Each tablet contains Erythromycin Stearate equivalent to Erythromycin 250mg.

Ettrocin for Suspension 200mg/5ml

When reconstituted each 5ml measure contains Erythromycin Ethylsuccinate equivalent to Erythromycin 200mg. Preservatives: Methylparaben 0.1% w/v and Propylparaben 0.01% w/v.

Pharmacological Information:

Erythromycin is usually bacteriostatic, but it may be bactericidal in high concentrations or against highly susceptible organisms. Erythromycin inhibits protein synthesis in susceptible organisms by binding to 50S ribosomal subunits, thereby inhibiting translocation of aminoacyl transfer-RNA and inhibiting polypeptides synthesis.

Erythromycin exerts its effect only against multiplying organisms. Only unionized erythromycin is believed to penetrate susceptible bacteria and penetration increases as the pKa of the drug is approached. Erythromycin generally penetrates the cell wall of gram-positive bacteria more readily than that of gram-negative bacteria and gram-positive organisms may accumulate 100 times more erythromycin than to gram-negative organisms.

Absorption of orally administered erythromycins occurs mainly in the duodenum. The bioavailability of the drugs is variable and depends on several factors including the particular erythromycin derivative, the formulation of the dosage form administered, acid stability, the presence of food in the GI tract and gastric emptying time.

Gastric acidity causes partial inactivation of some of these drugs, with the degree of inactivation depending on the acid stability of the particular derivative and dosage form. Erythromycin base is highly susceptible to gastric acid inactivation. Erythromycin ethylsuccinate is partially dissociated in the intestine. Both erythromycin and the undissociated ester are absorbed. In the blood, the ester is partially hydrolysed to release free erythromycin.

Single oral doses of the erythromycins generally produce peak serum concentrations within 1-4 hours. Higher peak serum concentrations are achieved when the drugs are given orally on 4 doses daily than following single doses. In general, oral administration 400 mg of erythromycin as the ethylsuccinate, 4 times daily maintains antibacterial serum concentrations of 0.1-2 ug/ml.

Erythromycin is widely distributed into most body tissue and fluids. All tissues except the brain have erythromycin concentrations which are higher and persist longer than serum concentration. Only low concentrations of Erythromycin (2-13% of serum concentrations) appear in the CSF.

The serum half-life of erythromycin in patients with normal renal function is usually 1.5-2 hours, but may range from 0.8-3 hours. In anuric patients, the half-life may be prolonged to 6 hours, but this is not considered to be clinically important. Although there are no reports on the half life of erythromycin in patients with impaired hepatic function, the possibility that the half-life may be prolonged should be considered.

Erythromycin is partly metabolized in the liver by N-demethylation to inactive, unidentified metabolites. Erythromycin is mainly excreted unchanged via the bile. Some reabsorption follows biliary excretion. Small amounts of erythromycin are also excreted in the urine.

Indication:

For the prophylaxis and treatment of infections caused by erythromycin-sensitive organisms. Clinical indications:

1. Upper respiratory tract infections
2. Lower respiratory tract infections
3. Skin and soft tissue infections
4. Bone infections
5. Sexually transmitted diseases
6. Oral/dental infections
7. Eye infections
8. Gastro intestinal infections
9. Prophylaxis

Microbiological indications: Erythromycin has been shown to be active in vitro against the following organisms. Staphylococci, Streptococci, Haemophilus influenzae L-forms, Mycoplasma pneumoniae, Legionella pneumophila, Branhamella, catarrhalis Bordetella pertussis, Corynebacterium diphtheriae (as an adjunct to antitoxin). Neisseria Treponema pallidum, Chlamydia trachomatis, Clostridia, Ureaplasma urealytica, Campylobacter.

Dosage and administration:

Administered orally.

Erythromycin preparations are taken 30 minutes or more before food.

Ettrocin 250 Film-Coated Tablet (Erythromycin Stearate):

Adults and children over 8 years:

For mild to moderate infections 1-2 daily in divided doses. For severe infections this may be increased to 4 daily in divided doses. Tablets should be taken before or with meals.

Elderly:

No special dosage recommendations.

Children under 8 years:

Erythromycin suspension is recommended.

Ettrocin for Suspension 200mg/5ml (Erythromycin Ethylsuccinate)

Adult:

400 mg four times daily or 800 mg twice daily.

Infants and children doses:

30 to 50 mg per kg of body weight daily in divided doses every 6 hours.

Contraindication:

Hypersensitivity to erythromycin.

Warning and Precautions:

Hepatic dysfunction - May occur, with or without jaundice, in patients receiving oral erythromycin.

Patients with hepatic impairment - Use with caution and watch for signs of excessive accumulation since drug is excreted primarily by the liver.

Concomitant theophylline therapy - May produce elevated serum theophylline, reduce theophylline dosage accordingly.

Suspected syphilitic lesions - If syphilis is suspected, perform a dark-field examination before instituting erythromycin stearate therapy and monthly serology tests for at least 4 months thereafter. Spinal fluid examinations should be done before treatment and as part of follow-up therapy.

Ototoxicity - Reversible hearing loss has occurred, primarily in patients who had renal impairment or received high doses.

Superinfection - Prolonged and repeated use may cause overgrowth of nonsusceptible bacteria or fungi, if superinfection occurs, discontinue use.

Interactions With Other Medicaments:

The use of erythromycin in patients who are receiving digoxin, warfarin, carbamazepine and high doses of theophylline may result in the potentiation of their effects due to reduction in the rates of excretion. Erythromycin may reduce the bactericidal action of lincomycin.

Pregnancy and Lactation:

Safety for use during pregnancy has not been established: There is no evidence of hazard from erythromycin in human pregnancy. It has been in wide use for many years without apparent ill consequence. Animal studies have shown no hazard. Erythromycin crosses the placental barrier, but fetal plasma levels are low.

Trace amount of erythromycin have been observed in human milk.