

Insert

CHLORTRALIM

OPHTHALMIC OINTMENT (Chlortetracycline HCl 1%)



NAME AND STRENGTH OF ACTIVE SUBSTANCES :

Chlortetracycline HCl 10 mg per 1g

PRODUCT DESCRIPTION :

Chlortralim Ophthalmic Ointment is a yellow homogenous sterile ointment in suitable ointment base for ophthalmic use, fill in aluminum ophthalmic ointment tube.

PHARMACODYNAMICS :

Chlortetracycline is a product of the metabolism of the organism *Streptomyces aureofaciens* and is one of the family of tetracycline antibiotics. Chlortetracycline is primarily bacteriostatic and is thought to exert its antimicrobial effect by the inhibition of protein synthesis. Chlortetracycline is effective against a wide range of Gram-positive and Gram-negative organism. The drugs in the tetracycline family have closely similar antimicrobial spectra, and cross resistance among them is common. Microorganism may be considered susceptible if the M.I.C (minimum inhibitory concentration) is not more than 4.0 mcg/ml and intermediate if the M.I.C is 4.0 to 12.5 mcg/ml. A tetracycline disc may be used to determine microbial susceptibility to drugs in the tetracycline family.

PHARMACOKINETICS :

Chlortetracycline, in an extremely fine form is incorporated in the ointment base. Absorption when applied to the eye without occlusion only results in slight absorption which is bounded to plasma protein, but is rapidly deactivated.

INDICATIONS :

Chlortralim Ophthalmic Ointment is indicated in eye infections caused by the following organism:

Richettsiae, *Mycoplasma pneumoniae*, agents of *psittacosis*, *ornithosis*, agents of *Lymphogranuloma venereum* and *Granuloma inguinale*, *Borrelia recurrentis*, *Haemophilus ducreyi*, *Pastuerella pestis*, and *Pastuerella tularensis*, *Bartonella bacilliformis*, *Bacteroides sp.*, *Vibrio comma* and *Vibrio fetus*, *Brucella sp.*, *E. coli*, *Enterobacter aerogenes*, *Shigella sp.*, *Haemophilus influenza*, *Klebsiella sp.*, *Streptococcus sp.*, *Diplococcus pneumoniae*, *Staphylococcus aureus*, *Neisseria gonorrhoeae*, *Treponema pallidum*, *Treponema pertenue*, *Listeria monocytogenes*, *Clostridium sp.*, *Bacillus anthracis*, *Actinomyces sp.*

RECOMMENDED DOSAGE :

Apply sparingly to the upper eye lid, close eye and move the eyeball about with the eye closed. The eye may then be covered with a light dressing. Repeat 4 to 6 times daily.

ROUTE OF ADMINISTRATION :

Instill into the conjunctival sac.

CONTRAINDICATION :

Chlortetracycline is contraindicated in patients with known hypersensitivity to the drug.

WARNINGS AND PRECAUTIONS :

Chlortralim should be applied sparingly to the affected area and it may cause some staining to the skin or clothing.

INTERACTION WITH OTHER MEDICAMENTS :

Not Available.

PREGNANCY AND LACTATION :

Not Available.

ADVERSE EFFECTS/ SIDE EFFECTS :

Hypersensitivity, skin rash, photosensitivity, local irritation at the site of application had been known to occur.

OVERDOSE AND TREATMENT :

Due to the low chlortetracycline content and its small pack, overdose is extremely rare. The symptoms are due to the ingestion of the petrolatum and is manifested by diarrhea or gastro-intestinal disturbance. Emetic or gastric lavage may be of value.

STORAGE CONDITION :

Store in a temperature below 30°C.
Protect from light.

PACKING AVAILABLE :

Tubes of 1/8 oz. (3.5 g).

MANUFACTURER :

ATLANTIC LABORATORIES CORPORATION LTD.

111 Moo 7 Bang Phli Noi, Bang Bo, Samut Prakan 10560, Thailand.

PRODUCT REGISTRATION HOLDER :

ATLANTIC LABORATORIES (M) SDN. BHD.

S-G-11, Gaya Resort Homes, Persiaran Kemuning Damai, Seksyen 32, 40460 Shah Alam, Selangor, Malaysia.

DATE OF REVISION :

03/12/2025

Atlantic Laboratories Corporation Ltd.
Samut Prakan, Thailand.

031225 MAL(R)