

ANTASONE Cream

[DESCRIPTION] Antasone cream is a white uniform cream.

[COMPOSITION] Each gram contains

Betamethasone valerate 1mg (Base)

Gentamicin sulfate 1mg (Potency)

Preservative: Methyl Paraben 1mg

Propyl Paraben 0.2mg

[PHARMACOLOGY/PHARMACOKINETICS]

Following topical application, corticosteroids produce antiinflammatory, antipruritic and vasoconstrictor actions, while Gentamicin is usually bactericidal in action.

Following topical application of a corticosteroid to most areas of normal skin, only minimal amounts of the drug reach the dermis and subsequently the systemic circulation; however, absorption is markedly increased when the skin has lost its keratin layer and can be increased by inflammation and/or diseases of the epidermal barrier (e.g. psoriasis, eczema).

Although the exact mechanism of action of Gentamicin has not been fully elucidated, the drug appears to inhibit protein synthesis in susceptible bacteria by irreversibly binding to 30S ribosomal subunits.

[INDICATIONS]

1. Allergic and inflamed dermatitis: acute eczema, chronic eczema, contact dermatitis, atopic dermatitis, seborrheic dermatitis, chronic lichen simplex, intertrigo, desquamative dermatitis, urticaria papulosa, psoriasis, pruritus itching (anus, pudenda), solar dermatitis.

2. It is used in the treatment of impetigo, erythema and other localised primary bacterial infection with a gram-negative component and also for secondary bacterial infections complicating other pre-existing dermatoses.

[DOSAGE AND ADMINISTRATION]

A small amount of Antasone cream should be applied to the lesion one to several times daily.

[CONTRAINDICATIONS]

Antasone cream is contraindicated as follows:

1. Tuberculosis of the skin, herpes simplex, varicella, herpes zoster, syphilis
2. Patients with a history of hypersensitivity to this drug.
3. Eczematoid external auditory meatusitis with drum membrane perforation.
4. Ulcer (Behcet's syndrome), burn of second-degree or more perforation.
5. Patients with a history of hypersensitivity to aminoglycosides antibiotics; streptomycin, kanamycin, gentamicin, neomycin etc. or bacitracin.

[PRECAUTIONS]

1. Pregnancy : Because the safety has not been established in pregnant women, prolonged use or large amounts of this drug should be avoided in the treatment of pregnant women.

2. Do not administer for ophthalmic use.

Use of topical Gentamicin preparation in closed hospital setting is actively discouraged.

[INCOMPATIBILITIES] None

[ADVERSE REACTIONS]

1. Skin infection : Mycotic infection (candidiasis, trichophytosis), bacterial infection (infectious impetigo, folliculitis) can occur. When these symptoms occur, especially in case of occlusive dressing, use together adequate antifungal agent or antibacterial agent and then the symptoms are not developed fast, discontinue use.
2. Other skin symptoms: In case of long-term administration, steroidal vulgaris(it is similar to acne vulgaris but induce white acnes), steroidal skin(skin atrophy, telangiectasia), steroidal rosacea namely dermatitis mouth surrounding(erythema, telangiectasia, crust, on mouth surrounding or occasionally full face), petechia, hypertrichosis, depigmentation may occur, when these symptoms occur, reduce amount and choose other drug without adrenal corticosteroids.
3. Hypersensitivity : When irritation or rash occur, discontinue use.
4. Pituitary, adrenal cortex: A large amount or long-term administration and occlusive dressing may inhibit the function of pituitary, and adrenal cortex.
5. Eye: In case of administration to eye lid can occur intra-ocular pressure acceleration, glaucoma. By a large amount or long term administration and occlusive dressing, cataract and glaucoma may occur.
6. Super infection : Occasionally infection of unsusceptive bacteria to this drug may occur, in this case, discontinue use.
7. Forbid long term administration because nephrotoxicity and dysacusis may occur.

[STORAGE] Store in a well-closed container at 30°C or below.

[EXPIRY] 3 years.

[PACK SIZE] 10g tube & 20x10g tube

MANUFACTURER:

SHIN POONG PHARM CO LTD

7,Wonsi-ro, Danwon-gu,Ansan-si,
Gyeonggi-do, Republic of Korea.

PRODUCT REGISTRATION HOLDER:

The Zyfas Medical Co.,

No. 7, Jalan Molek 3/10,
81100 Johor Bahru.

Revised Date: March 2022