

[DUOPHARMA (M) SDN BHD]

**GENTA – DEX EYE DROPS
FOR OPHTHALMIC USE ONLY**

DESCRIPTION:

Genta-Dex Eye Drops is a clear, almost colourless liquid.

COMPOSITION:

Each ml of Genta-Dex Eye Drops contains:

Gentamicin Sulphate equivalent to Gentamicin 0.3% w/v

Dexamethasone Sodium Phosphate equivalent to Dexamethasone Phosphate 0.1% w/v

Preservative used: Benzalkonium Chloride 0.01% w/v

PHARMACODYNAMICS:

Ophthalmic eye drops combines the potent anti-inflammatory and anti-allergic action of Dexamethasone Sodium Phosphate with the broad spectrum bactericidal activity of the aminoglycoside gentamicin sulphate. In vivo, staphylococcal species have responded to gentamicin sulphate. In vitro, gentamicin sulphate is active against a wide variety of pathogenic Gram-negative and some Gram-positive bacteria: a coagulase-positive and coagulase-negative staphylococci, including certain strains that are resistant to penicillin. *Escherichia coli*, *Proteus* species (indole-positive and indole-negative), *Pseudomonas aeruginosa*, species of the *Klebsiella-Enterobacter-Serratia* group, *Haemophilus influenzae*, species of *Citrobacter*, *Salmonella*, *Shigella*, *Moraxella*, *Serratia* and *Neisseria*, particularly the gonococcus. Gentamicin eye drops did not produce an effective concentration in the aqueous humour.

PHARMACOKINETICS:

Gentamicin sulfate is a water soluble antibiotic of the aminoglycoside group which has shown activity against a wide variety of pathogenic gram-negative and gram-positive bacteria. The gram-positive bacteria against which gentamicin sulfate is active include coagulase positive and negative staphylococci.

Dexamethasone is readily absorbed from the gastrointestinal tract. Its biological half-life in plasma is about 190 minutes. Binding of dexamethasone to plasma proteins is less than for most other corticosteroids. Up to 65% of a dose is excreted in urine within 24 hours. Clearance in premature neonates is reported to be proportional to gestational age, with a reduced elimination rate in most premature. It crosses the placenta.

INDICATIONS:

Genta-Dex eye drops is indicated for the treatment of allergic and inflammatory conditions of the eye where there is superficial bacterial infection of the conjunctiva, cornea (without epithelial defect) and margin of eye lid. This includes conjunctivitis, keratitis and keratoconjunctivitis, corneal ulcers, blepharitis and blepharoconjunctivitis, acute meibomianitis and dacryocystitis.

NB: "To ensure that fungal infection has been eliminated before use of preparation."

RECOMMENDED DOSAGE:

Instill two drops of Genta-Dex Eye Drops three to four times daily. As with eye medication, Genta-Dex Eye Drops may sting briefly on application.

ROUTE OF ADMINISTRATION: For ophthalmic use

CONTRAINDICATIONS:

Genta-Dex Eye drops is contraindicated in epithelial herpes simplex keratitis (dendritic keratitis), vaccinia, varicella, and viral diseases of the cornea and conjunctiva, mycobacterial or fungal infections of the eye, trachoma, or hypersensitivity to any component of this preparation.

Use of corticosteroid / antibiotic combinations is contraindicated after removal of a corneal foreign body.

WARNING & PRECAUTIONS:

If prompt clinical response is not obtained with the use of Genta-Dex Eye Drops, further evaluation is advised. When Genta-Dex Eye Drops is applied to the eye for 10 or more days, tonometry and slit lamp examination are advisable. In those diseases causing thinning of the cornea or sclera, perforation has been known to occur with the use of topical corticosteroids.

[DUOPHARMA (M) SDN BHD]

Accordingly, it is not advisable to treat bacterial corneal ulcers which may be due to *Pseudomonas aeruginosa*, with a combination antibiotic-antiinflammatory product as initial therapy. It is prudent to use an anti-infective agent alone initially. For ulcers caused by addition of an anti-inflammatory agent to minimize the fibrous reaction and scarring of the cornea is suggested. In acute purulent of the eye, corticosteroids may mask infection or enhance existing infection. Prolonged use of topical antibiotics or corticosteroids may give rise to overgrowth of non-susceptible microorganisms and fungi. Should this occur, or if irritation or hypersensitivity to Genta-Dex Eye Drops develops, discontinue use of this preparation and institute appropriate therapy. Cross allergenicity among aminoglycosides or corticosteroids has been demonstrated. Contamination of the eye drops may occur if the dispenser tip touches any surface. Safety and effectiveness of Genta-Dex Eye Drops in children below the age of eight years have not been established.

Use of topical gentamicin preparations in closed hospital settings is actively discouraged.

INTERACTION WITH OTHER MEDICAMENTS:

Co-treatment with CYP3A inhibitors, including cobicistat-containing products, is expected to increase the risk of systemic side-effects. The combination should be avoided unless the benefit outweighs the increased risk of systemic corticosteroid side-effects, in which case patients should be monitored for systemic corticosteroid side-effects.

USE IN PREGNANCY & LACTATION:

Use in Pregnancy: Gentamicin and other aminoglycosides cross the placenta. There is evidence of selective uptake of gentamicin by the foetal kidney resulting in damage (probably reversible) to immature nephrons. Eighth cranial nerve damage has also been reported following in-utero exposure to some of the aminoglycosides. Because of their chemical similarity, all aminoglycosides must be considered potentially nephrotoxic and ototoxic to the foetus. It should also be noted that therapeutic blood levels in the mother do not equate with safety for the foetus.

Use in Lactation: Since it is not known whether the components of Genta-Dex Eye Drops are excreted in human milk, caution should be exercised when it is administered to a nursing woman.

SIDE EFFECTS:

Adverse effects due to ophthalmic corticosteroids include increased intraocular pressure; glaucoma; infrequent optic nerve damage; defects in visual acuity and visual fields; posterior subcapsular cataract formation; delayed wound healing; acute anterior uveitis; perforation of the globe; mydriasis; loss of accommodation; ptosis. Transient eye irritation may occur with ophthalmic gentamicin sulphate. The extensive use of steroids may cause systemic side effects and ocular herpes simplex has occurred in patients under systemic or local corticosteroid therapy for other conditions. Extended use of topical steroids may cause increased intraocular pressure in certain individuals. It is advisable that intraocular pressure be checked frequently. In those diseases causing thinning of the cornea, perforation has been known to occur with the use of topical steroids.

Less frequent incidence: Itching; redness, swelling, burning and stinging.

SYMPTOMS AND TREATMENT OF OVERDOSE:

In case of overdosage, immediately flush the eye(s) with water or normal saline.

STORAGE CONDITIONS:

Store below 25°C. Protect from light. Do not use one month after first opening.
Keep out of reach of children. *Jauhkan daripada kanak-kanak.*

PACK SIZE:

Pack 1 bottle of 5ml in a 10ml eye drop bottle per box.

SHELF LIFE:

Please refer to outer package .

PRODUCT REGISTRATION HOLDER & MANUFACTURER:

DUOPHARMA (M) SDN BHD

Lot 2599, Jalan Seruling 59 Kawasan 3,
Taman Klang Jaya, 41200 Klang,
Selangor, Malaysia.

15000