

DYNA AMPICILLIN

	DESCRIPTION	CONTENT
DYNA AMPICILLIN CAPSULE 250MG MAL19910018AZ	CAPSULE Capsule Colour : Red/Black Capsule Size : # 2 Granules Colour : White	Each capsule contains: Ampicillin Trihydrate equivalent to Ampicillin..... 250 mg
DYNA AMPICILLIN CAPSULE 500MG MAL19910019AZ	CAPSULE Capsule Colour : Red/Black Capsule Size : # 0 Granules Colour : White	Each capsule contains: Ampicillin Trihydrate equivalent to Ampicillin 500 mg

PHARMACODYNAMICS:

Ampicillin is a bactericidal antibiotic which inhibits bacterial cell wall synthesis, probably by acylation of membrane-bound transpeptidase enzymes. This prevents cross-linkage of peptidoglycan chains which is necessary for bacterial cell wall strength and rigidity. Also, cell division and growth are inhibited and lysis of susceptible bacteria frequently occurs. Rapidly dividing bacteria are most susceptible to penicillin group of antibiotics including Ampicillin.

PHARMACOKINETICS:

Ampicillin is acid stable and 30 to 60% of an oral dose is absorbed. Peak serum concentrations of 0.8 to 4 mcg/ml are attained 1 to 2 hours after oral doses of 125 to 500 mg of anhydrate or trihydrate. The serum half-life is 1 to 2 hours which is increased in subjects with renal function impairment. Ampicillin enters most body cavities, tissues and fluids except the cerebrospinal fluid unless the meninges are inflamed, in which case large amounts may be detectable in this fluid; Ampicillin crosses the placenta and is excreted in the milk; volume of distribution about 20 L which in subjects with cirrhosis may reach about 60 L. About 20% of Ampicillin are bound to plasma proteins. 30% to 50% of an oral dose is excreted in the urine in 6 to 8 hours; up to 70% is excreted in 24 hours; urinary excretion is decreased after the administration of Probenecid; Ampicillin is excreted in the bile in concentrations up to 4 mcg/ml after an oral dose of 125 mg; biliary excretion is increased in subjects with cirrhosis.

INDICATIONS:

Infections caused by sensitive gram-positive and gram-negative microorganism.

ENT infections: Otitis media, sinusitis, tonsillitis, pharyngitis, laryngitis, epiglottitis.

Lower respiratory tract infections: Acute and chronic bronchitis, pneumonia, pleuritis.

Urinary tract infections: Acute and recurrent cystitis, acute and chronic pyelonephritis, asymptomatic bacteriuria, prostatitis.

Skin and soft tissue infections: Erysipelas, impetigo, pyoderma.

RECOMMENDED DOSE:

Adults: 250 - 500 mg every 6 hours up to 6 g daily.

Children up to 20 kg body-weight: 12.5 - 25 mg per kg body-weight every 6 hours.

Gonorrhoea: 3 g as a single oral dose may be administered with 1 g Probenecid.

Gastro-intestinal infections due to sensitive organisms: 1 - 2 g orally every 6 hours.

Note :

- * Patients should continue medicine for full course of treatment.
- * Take medicine on empty stomach.
- * Reduced dosage is required in patients with impaired renal function.

ROUTE OF ADMINISTRATION:

FOR ORAL USE ONLY

CONTRAINDICATIONS:

Patients hypersensitive to Ampicillin.

WARNINGS AND PRECAUTIONS:

Patients hypersensitive to other penicillins or cephalosporins or penicillamine may be hypersensitive to Ampicillin.

Reduced dose may be required in patients with impaired renal function. It decreases the efficacy of oestrogen containing oral contraceptives.

It may also affect the absorption of other drugs due to its effect on the gastrointestinal flora.

Serious and occasionally fatal hypersensitivity reactions (including anaphylactoid and severe cutaneous adverse reactions) have been reported in patients on penicillin therapy with beta-lactams. Before initiating therapy with Dyna Ampicillin Capsule, careful inquiry should be made concerning previous hypersensitivity reactions to penicillins, cephalosporins, carbapenems or other beta-lactam agents. If an allergic reaction occurs, Dyna Ampicillin Capsule must be discontinued immediately and appropriate alternative therapy instituted.

DRUG INTERACTIONS:

Bacteriostatic drugs may interfere with the bactericidal action of Ampicillin.

In common with other oral broad-spectrum antibiotics, Ampicillin may reduce the efficacy of oral contraceptives and patients should be warned accordingly.

Probenecid decreases the renal tubular secretion of Ampicillin. Concurrent use with Ampicillin may result in increased and prolonged blood levels of Ampicillin.

Concurrent administration of Allopurinol during treatment with Ampicillin can increase the likelihood of allergic skin reactions.

It is recommended that when testing for the presence of Glucose in urine during Ampicillin treatment, enzymatic glucose oxidase methods should be used. Due to the high urinary concentrations of Ampicillin, false positive readings are common with chemical methods.

PREGNANCY AND LACTATION:

Risk-benefit must be considered when given to pregnant women and during breast-feeding.

SIDE EFFECTS:

Hypersensitivity or allergic reactions i.e. skin rashes either urticarial or maculopapular, itching or wheezing.

GIT irritation i.e. abdominal or stomach cramps, pain, bloating, nausea, vomiting, diarrhoea & pseudomembranous colitis.

Fever, increased thirst, unusual tiredness or weakness.

Supra-infection with *Pseudomonas* & *Candida* with prolonged use.

Skin and subcutaneous tissue disorders:

Frequency 'very rare': Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS).

SYMPTOMS AND TREATMENT OF OVERDOSE:

Overdose may cause GIT disturbances, reversible impairment of liver function and diarrhoea. Treatment includes stop antibiotic treatment, treat GIT distress by giving milk every 3 hours alternating with Bismuth Subcarbonate 5 g every 3 hours.

Problems of overdose are unlikely to be encountered. Gross overdose will produce very high urinary concentration, more so after parenteral administration. Problems are unlikely if adequate fluid intake and urinary output are maintained; however crystalluria is a possibility. More specific measures may be necessary in patients with impaired renal function; the antibiotic is removed by haemodialysis.

PACKING/PACK SIZE(S):

DYNA AMPICILLIN CAPSULE 250MG - Blister pack of 100x10's.

DYNA AMPICILLIN CAPSULE 500MG - Blister pack of 50x10's.

JAUHI UBAT DARIPADA KANAK-KANAK
KEEP OUT OF REACH OF CHILDREN

MANUFACTURER/PRODUCT REGISTRATION HOLDER:

DYNAPHARM (M) SDN. BHD. 198001011897 (65683-V)

2497, Mk. 1, Lorong Perusahaan Baru 5,
Kawasan Perusahaan Perai 3,
13600 Perai, Pulau Pinang, MALAYSIA

PI1PC D020-01
Latest Revision: Jul 2023