

ROYCE®

Mebendazole Tablet 100mg (Mebendazole-Anthelmintics)

Presentation:

A cream coloured, round, plain, shallow biconvex tablet with a diameter of 10.3 mm.

Content:

Each tablet contains:
Mebendazole 100mg

Pharmacological Information:

The anthelmintic action of Mebendazole was considered to be due to its inhibition of glucose uptake by nematodes thereby depleting the energy level until it becomes inadequate for survival.

Mebendazole is not significantly absorbed from the gastrointestinal tract only about 2% of a dose is excreted unchanged or as a metabolite in the urine.

Indication:

Mebendazole is an anthelmintic effective against:

Enterobiasis (Threadworm infection)

Ascariasis (roundworm infection)

Trichuriasis (whipworm infection)

Uncinariasis (hookworm infection) caused by *Necator americanus* and *Ancylostoma duodenale*.

Dosage and Administration:

For threadworm infection:

Adult: 100 mg given in a single dose; if patient is not cured 3 weeks later, repeat dosage.

Child (over 2 years): same as adult.

For hookworm and whipworm infections:

Adult: 100 mg twice daily (morning and evening) for 3 consecutive days; if patient is not cured 3 weeks later, repeat dosage.

Child (over 2 years): same as adult.

Safety for use in children under 2 years of age has not been established.

Contraindication:

Mebendazole is contraindicated in pregnant women and in persons who have shown hypersensitivity to the drug.

Warning and Precautions:

Pregnancy

Caution patients about the potential risk of use during pregnancy (see USE IN PREGNANT AND LACTATION).

Cleanliness

Inform patients that cleanliness is important in preventing transmission of disease and reinfection. Hydatid disease. There is no evidence that Mebendazole is effective, even at high doses, in the treatment of hydatid disease.

Interactions with Other Medicaments

No clinically significant drug interactions have been identified.

Pregnancy and Lactation:

Embryotoxic and teratogenic effects have been detected in rats. Use during pregnancy, particularly during the first trimester, is not recommended, administer only if the expected benefits justify the potential risk to the fetus. It is now known whether Mebendazole is excreted in human milk; use with caution in nursing mothers.

Should not be administered during confirmed or suspected pregnancy.

Side Effects:

Gastrointestinal - Transient abdominal pain and diarrhoea in cases of massive infection and expulsion of worms.

Symptoms and Treatment of Overdose:

Sign and symptoms - Gastrointestinal reactions persisting for up to a few hours may occur.

Treatment - Induce vomiting and purging.

Storage Condition:

Store in a dry place (below 30°C). Keep container tightly closed. Protect from light.

Pack Size:

Plastic bottle of 30 tablets.

Product Registration Number:

MAL19910328XZ

Further information can be obtained from your doctor or pharmacist.



Product Holder / Manufactured by:

ROYCE PHARMA MFG. SDN.BHD. (650435-X)

PT 1663, Nilai Industrial Estate,

71800 Nilai, Negeri Sembilan, Malaysia.

Revision date 201217