



PARACETAMOL-250 SUSPENSION (ORANGE)

COMPOSITION:

Each 5ml contains
Paracetamol 250mg

Methylparaben 0.1% w/v as preservative.
Sunset Yellow Dye 0.015% w/v as colourant.

PRESENTATION:

An orange coloured and orange flavoured suspension.

INDICATIONS:

For the relief of fever, symptoms of cold and influenza, teething pain and headache.

DOSAGE AND ADMINISTRATION:

For oral administration only.

SHAKE WELL BEFORE USE.

Children

1-6 years old : 2.5 - 5ml.

7-12 years old : 5 - 10ml.

To be taken 3 or 4 times a day.

Children 3 months to 1 year are advised to use Axcel Paracetamol-120 Syrup.

CONTRAINDICATION:

Contraindicated in patients with nephropathy.

PRECAUTIONS:

It should be given with care to patients with impaired kidney or liver functions. Chronic use should be avoided.

THIS PREPARATION CONTAINS PARACETAMOL.
DO NOT TAKE ANY OTHER PARACETAMOL.
CONTAINING MEDICINES AT THE SAME TIME.

Allergy alert : Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

SIDE EFFECTS:

Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome / Toxic Epiderma Necrolysis have been reported.

DRUG INTERACTION:

The prothrombinopenic effect of the oral anticoagulants may be increased by chronic

administration of full doses of paracetamol.

OVERDOSAGE & TREATMENT:

In non-fatal cases, the hepatic lesions are reversible over a period of weeks or months. Measurement of the plasma half-life of paracetamol during the first day of acute poisoning provides an early indication of the severity of the liver injury. Hepatic necrosis should be anticipated if the half-time exceeds 4 hours, and hepatic coma is likely if the half-time is greater than 12 hours. Encephalopathy should also be anticipated if serum bilirubin concentration exceeds 4mg / 100ml during the first 5 days. Treatment of acute paracetamol overdose is purely symptomatic, vigorous supportive therapy is essential in severe intoxication. Induction of vomiting or gastric lavage should be performed in all cases, follow by oral administration of activated charcoal. Hemodialysis, if it can be initiated within the first 12 hours, may be advocated for patient with a plasma paracetamol concentration greater than 120mg / ml 4 hours after drug ingestion. Peritoneal dialysis with albumin in the dialysate may be useful. Cysteamine of N-acetyl-cysteine intravenously, or 1-methionine orally appear to protect the liver if administered within 10 hours of ingestion and overdose. There are no known antidotes of paracetamol.

STORAGE:

Keep container well closed. Store below 30°C. Protect from light.

**KEEP OUT OF REACH OF CHILDREN
JAUHI DARI KANAK-KANAK****PACK QUANTITIES:**

Available in PET bottle of 60ml, 90ml and 100ml and glass bottle of 90ml & 120ml.

Further information can be obtained from doctor, pharmacist or the manufacturer.

MAL19911435XZ

Manufactured by & Product Registration Holder :



Kotra Pharma (M) Sdn. Bhd.
No.1, 2 & 3, Jalan TTC 12, (90082-V)
Cheng Industrial Estate,
75250 Melaka, Malaysia.

REGN-151122
Date of revision : 15/11/22