



PRITAMOL SUSPENSION 250MG/5ML

Description: Pink, raspberry flavoured suspension.

Contents:

Pritamol Suspension : Paracetamol 250mg
Methylparaben 0.1% w/v
Propylparaben 0.01% w/v

This preparation contains PARACETAMOL.
Do not take any other PARACETAMOL
containing medicines at the same time.

Indications:

For the relief of fever and pain.

Pharmacology:

Paracetamol, an analgesic and antipyretic, acts by inhibiting prostaglandin synthesis in the CNS and, to a lesser extent, through a peripheral action by blocking pain-impulse generation. It produces antipyresis by acting centrally on the hypothalamic heat-regulating center to produce peripheral vasodilation resulting in increased blood flow through the skin, sweating, and heat loss. Oral absorption is rapid and elimination is via hepatic metabolism and renal excretion.

Adverse Effects / Undesirable Effect:

In recommended therapeutic dosage, paracetamol is usually well tolerated. Rarely, hypersensitivity and hematological reactions occur.

Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome/ Toxic Epidermal Necrolysis have been reported.

Warnings and Precautions:

- Caution in patients with hepatic or renal function impairment, and in patients regularly taking other hepatic enzyme inducing agents.
- Chronic use should be avoided, unless otherwise directed by physician.

Allergy Alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Overdosage:

Clinical features include diarrhoea, loss of appetite, nausea or vomiting, stomach cramps, unusual increase in sweating, and hepatic or renal failure.

Treatment consists of emesis or gastric lavage, if appropriate. Oral administration of acetylcysteine should be instituted as soon as possible or within 10 to 12 hours after ingestion of an overdose. The recommended adult dose of acetylcysteine is 140 mg/kg body weight initially, then 70 mg/kg body weight every 4 hours for 17 doses. Any dose vomited within 1 hour of administration must be repeated.

Dosage and administration: (ORAL)

Adult 2-4 teaspoonfuls, 3-4 times daily.

Children 1-6 years, 1/2-1 teaspoonful 3-4 times daily.

Children 7-12 years, 1-2 teaspoonfuls 3-4 times daily.

Storage : Keep container tightly closed. Store in a cool, dry place away from light, below 30°C
Keep out of reach of children.

Presentation/Packing : Suspension 250mg/5ml x 60ml

Shelf life : 2 years

Manufacturing & : **PRIME PHARMACEUTICAL SDN. BHD.**

Distributed by : 1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah
14000 Bukit Mertajam, Penang, Malaysia.