



PRIMEDAZOL TABLET 100mg, 500mg
PRIMEDAZOL SUSPENSION 100mg/5ml

Contents:

Primedazol Tablet 100mg : Mebendazole USP 100mg / tablet (Light orange, round with top-scored tablet)
Primedazol Tablet 500mg : Mebendazole USP 500mg / tablet (Yellow, round with top-scored tablet)
Primedazol Suspension : Mebendazole USP 100mg / 5ml (Light yellow, vanilla flavoured suspension)
Methylparaben 0.1% w/v
Propylparaben 0.01% w/v

Indications:

For treatment of single or mixed infestations of pinworm / threadworm (*Enterobius*), roundworm (*Ascaris*), whipworm (*Trichuris*), hookworm (*Ancylostoma* and *Necator*) and tapeworm.

Pharmacology:

Mebendazole, a broad-spectrum anthelmintic, blocks the glucose uptake by susceptible helminths, thereby depleting glycogen stored within the parasite. This results in reduced formation of adenosine triphosphate (ATP) required for survival and reproduction of the helminth. Oral absorption is insignificant and elimination is mainly by fecal excretion.

Adverse Effects:

Side effects are usually mild and transient; abdominal pain and diarrhoea may occur in cases of massive infection and expulsion of worms.

Precautions:

- No special preparations (e.g. dietary restrictions or fasting, purging, or cleansing enemas) are required before, during, or immediately after therapy.
- For patients on high-dose therapy, take with meals, especially fatty ones, to increase absorption.
- Safety for use in lactation has not been established.

Contraindications:

- It is not recommended during pregnancy.
- Avoid in patients who have shown hypersensitivity to the drug.

Dosage and Administrations:

Adults and children over 2 years:

For pinworm / threadworm infestations - Oral, 100mg as a single dose. Repeat if necessary after 2 to 3 weeks.

For other infestations - Oral, 100mg two times a day, morning and evening for three days.
or
Oral, 500mg as a single dose.
If necessary, a second course of treatment may be repeated in two to three weeks.

Note: The tablets may be swallowed whole or crushed and mixed with food.
Continue medication for full course of treatment.

Overdosage:

If accidentally overdose, gastrointestinal symptoms may occur and may last up to a few hours. Vomiting and purging should be induced.

Storage : Keep container tightly closed. Store in a cool, dry place below 30°C.
Protect from light. Keep away from children.

Shelflife : 3 years

Presentation/Packing : Tablet 100mg x 500's, 500mg x 100's, 100mg/5ml x 30ml

Manufactured & : PRIME PHARMACEUTICAL SDN.BHD.

Distributed by 1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah,
14000 Bukit Mertajam, Penang, Malaysia.