

DHAMOL

PRESENTATION:

DHAMOL SUSPENSION : Paracetamol BP (fine powder) 250 mg/5 ml

PRODUCT DESCRIPTION:

Homogenous pink suspension with strawberry flavour and sweet taste.

PHARMACOLOGY:

Paracetamol has analgesic and antipyretic effects. It inhibits the prostaglandin synthesis in the brain to relieve pains of moderate intensity, such as headache, dysmenorrhoea, pain in muscle and nerve disorders. It reduces fever by a direct effect on the heat-regulating centre to increase dissipation of body heat.

PHARMACOKINETIC:

It is rapidly and completely absorbed from the G.I.T. and metabolised in the liver. The hydroxylated metabolites are responsible for methemoglobin formation and hepatotoxicity. About 3% of the absorbed paracetamol is excreted unchanged in the urine and about 80% in conjugated forms.

INDICATIONS:

For relief of fever and pain of moderate intensity e.g. headache, dysmenorrhoea and toothache.

RECOMMENDED DOSAGE: Orally

Children:

Under 3 months : 5-10 mg/kg of body weight
or on doctor's advice

3 months - 1 year : 1.25 - 2.5 ml,

1 - 6 years : 2.5 - 5 ml,

6 - 12 years : 5 - 10 ml.

These doses may be given 4 to 6 hourly when necessary. (Maximum: Not more than 4 doses daily).

ADVERSE EFFECT/UNDESIRABLE EFFECT:

Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome/Toxic Epidermal Necrolysis have been reported.

Side effects of paracetamol are usually mild; skin rashes and allergic reactions occur occasionally.

PRECAUTIONS/WARNINGS:

This preparation contains PARACETAMOL. Do not take any other PARACETAMOL containing medicines at the same time.

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Inappropriate use of product containing paracetamol can result in liver toxicity or severe liver damage. Do not exceed the stated dose. Paracetamol should be given with care to patients with impaired kidney or liver functions. Chronic use should be avoided. Daily use may potentiate oral anticoagulants.

USE IN PREGNANCY AND LACTATION:

Not known to be harmful.

CONTRAINDICATION:

Nephropathy.

DRUG INTERACTION:

The absorption of paracetamol may be accelerated by metoclopramide. Excretion may be affected and plasma concentrations altered when administered with probenecid.

INCOMPATIBILITY:

Not reported.

OVERDOSAGE & TREATMENT:

Symptoms of overdosage in the first 24 hours are pallor, nausea, vomiting, anorexia and abdominal pain. Liver damage may become apparent 12 to 48 hours after ingestion. Consult a doctor immediately. Gastric lavage is carried out if the overdose was taken within 2 hours of admission. Antidotes such as acetyl cysteine and methionine may be necessary.

AVAILABILITY:

Suspension : 60ml in plastic container, 100ml in amber glass bottle.

STORAGE:

Keep out of the reach of children. Store below 30°C. Protect from light.

EXPIRY PERIOD:

Do not use after the expiry date stated on the label.

DATE OF REVISION:

November 2021

For further information, please consult your physician or pharmacist.

Manufactured by:
PT Actavis Indonesia
Jl. Raya Bogor Km. 28, Kelurahan Pekayon,
Kecamatan Pasar Rebo
Kota Jakarta Timur, Provinsi DKI Jakarta,
Indonesia

W
Actavis
AAAN8921
40958-02

Artwork

 PT. Gaya Serasi Indonesia	
Date	
From	
Product Name	Leaflet Dhamol Suspension 250 mg-MY
Code	40958-02
Colour	K
Material	HVS 70 gram
Finishing	Lipat
Size	125 x 220 mm

Quality Control	Manager Prod.	Marketing
Date :	Date :	Date :

Customer Approval Design & Colour	
Date :	

You Dream it We Print it