

WINMOTAN® SYRUP PARACETAMOL 250MG/5ML
(Analgesic - Antipyretic)

Active Ingredient:
Each 5 ml contains -
Paracetamol 250mg
Preservative -
Sodium Benzoate 0.1%w/v

Presentations:
Bottle of 30 ml.

Product Descriptions:
Clear colourless to light brown syrupy solution.

Pharmacology:
Paracetamol is an effective analgesic and antipyretic but lacks anti-inflammatory action. It indicated for symptomatic relief of mild to moderate pain and to reduce fever, especially in Aspirin sensitive patients and children. The mechanism of analgesic action has not been fully determined. Paracetamol may act by inhibiting prostaglandin synthesis in the central nervous system and through a peripheral action by blocking pain-impulse generation. The peripheral action may also be due to inhibition of the synthesis of prostaglandins or to inhibition of the synthesis or actions of other substances, which sensitize pain receptors to mechanical or chemical stimulation. Paracetamol probably produces antipyresis by acting centrally on the hypothalamic heat regulating centre to produce peripheral vasodilation resulting in increased blood flow through the skin, sweating and heat loss. The central action probably involves inhibition of prostaglandin synthesis in the hypothalamus.

Paracetamol readily absorbed following oral administration. Peak plasma concentration occurring about 30 minutes to 2 hours after ingestion. Metabolised in the liver and excreted in the urine mainly as glucuronide and sulphate conjugates. Less than 5% is excreted as unchanged Paracetamol. The elimination half life varies from about one to four hours. Plasma protein binding is negligible at usual therapeutic concentration but increase with increasing concentrations. Volume of distribution: About 1L/kg. Plasma clearance: About 5ml/ min/kg. Duration of action: 3 to 4 hours.

Indications:
For fever and relief of mild to moderate pain including headache and toothache.

Dosage and Administration: For oral administration only.

Adults and children above 12 years:

10ml to 20ml (2 to 4 teaspoonfuls).

Children 7 to 12 years:

5ml to 10ml (1 to 2 teaspoonfuls).

Children 1 to 6 years:

2.5ml to 5ml (1/2 to 1 teaspoon).

To be taken 3 to 4 times a day together with water.

Note: Administration to infants under 1 month is not recommended.

Symptoms and Treatments of Overdose:

Symptoms: Symptoms of Paracetamol overdosage in the first 24 hours are pallor, nausea, vomiting, anorexia and abdominal pain. Liver damage may become apparent 12 to 48 hours after ingestion by increases in serum concentrations of aminotransferases and bilirubin and in prothrombin time. Abnormalities of glucose metabolism and metabolic acidosis may occur. In severe poisoning, hepatic failure may progress to encephalopathy, coma and death. Acute renal failure with acute tubular necrosis may develop even in the absence of severe liver damage. Cardiac arrhythmias have been reported.

Treatment: Prompt specific treatment is essential:

Removal of stomach contents by aspiration and lavage.

Administration of activated charcoal should be considered.

Dextrose and blood infusion.

Use antidote.

Haemoperfusion may be worthwhile if too much time has elapsed since the poisoning to allow use of antidote.

Antidote: Cysteamine is sometimes used by injection but acetylcysteine or methionine is now preferred. Acetylcysteine is given by intravenous infusion in an initial dose of 150mg per kg body weight over 15 minutes followed by 50mg per kg over four hours and then 100mg per kg over the next 16 hours. Alternatively methionine 2.5g may be given by mouth every 4 hours to total of four doses.

Contraindications:

Patients with known hypersensitivity to Paracetamol.

Not recommended for babies under 1 month old.

Precautions:

This preparation contains Sodium Metabisulphite that may cause serious allergic type reaction in certain susceptible patients. Do not use if known to be hypersensitive to bisulphites.

Do not exceed the recommended dosage and continual use should not exceed 5 days at a time (10 days for adult) without physician guidance.

Use with caution in patients taking medication affecting liver, impaired kidney and liver functions.

This preparation contains Paracetamol. Do not take any other Paracetamol containing medicines at the same time.

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Use in Pregnancy and Lactation:

Category A: Paracetamol has been taken by large number of pregnant women and women of childbearing age without any proven increase in frequency of malformations or other direct or indirect harmful effects on the foetus having been observed.

Paracetamol is excreted in breast milk in less than 1% of the dose ingested by lactating women. Therefore, maternal ingestion of therapeutic doses does not present a risk to the baby.

Interactions With Other Medicaments:

Anticoagulant dosage may require reduction if Paracetamol medication is prolonged.

Undesirable Effects:

Side effects of Paracetamol are usually mild, though haematological reactions have been reported. Skin rashes and other allergic reactions occur occasionally.

Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome / Toxic Epidermal Necrolysis have been reported.

Storage:

Store below 30°C. Protect from light. Keep cap tightly closed.

Keep out of reach of children.

Jauhkan daripada capaian kanak-kanak.

Shelf Life:

3 years from the date of manufacture.

Manufacturer and Product Registration Holder

WINWA MEDICAL SDN. BHD.

199101099271(75734-A)

1952, Lorong I.K.S. Bukit Minyak 3,

Taman I.K.S. Bukit Minyak,

14000 Bukit Mertajam,

Pulau Pinang, Malaysia.

MAL19912819XZ

Date of Revision: 24 June 2025

X.LWM-I.13

Size: 75mm x 135mm
■ Dark Blue (Pantone 2767U)

Checked by	
Approved by	