

EROGRAN GRANULES

VIERO06-var (MY)

DESCRIPTION

White, cherry flavoured granules.

COMPOSITION

Erythromycin ethylsuccinate equivalent to 200 mg of erythromycin in 5 mL when reconstituted.

PHARMACODYNAMICS

Erythromycin, a bacteriostatic antibiotic, may be bactericidal in high concentrations or when used against highly susceptible organisms. It is thought to penetrate the bacterial cell membrane and to reversibly bind to the bacterial ribosomes so that translocation of peptides is prevented and subsequent protein synthesis is inhibited.

PHARMACOKINETICS

Oral absorption as the free erythromycin is reduced by food. Erythromycin is widely distributed throughout body tissues and fluids; it does not cross the blood brain barrier. Highest concentrations are found in the liver, bile and spleen. It is excreted in the bile and urine.

INDICATIONS

- Indicated in the treatment or prophylaxis of the following:
- Upper and lower respiratory tract infections.
 - Skin and soft tissue infections including erythrasma.
 - Genitourinary tract infections including gonorrhoea and syphilis.
 - Intestinal amoebiasis.
 - Others include bacterial endocarditis, streptococcal pharyngitis, diphtheria, Legionnaires' disease and pertussis.

CONTRAINDICATIONS

Avoid in patients known to be hypersensitive to erythromycins.

WARNING AND PRECAUTIONS

- Caution in patients with impaired hepatic function.
- Safety for use in pregnancy and lactation has not been established.
- Following prolonged or repeated erythromycin therapy, overgrowth of non susceptible bacteria or fungi may occur.
- Should not be used in hypersensitive patients.
- Rare cases of serious cardiovascular adverse event including deaths, cardiac arrests, torsades de pointes and other ventricular arrhythmia have been observed, when used in patients taking concomitant terfenadine or astemizole.
- There have been reports of infantile hypertrophic pyloric stenosis (IHPS) occurring in infants following erythromycin therapy. In one cohort of 157 newborns who were given erythromycin for pertussis prophylaxis, seven neonates (5%) developed symptoms of non-bilious vomiting or irritability with feeding and were subsequently diagnosed as having IHPS requiring surgical pyloromyotomy. Since erythromycin may be used in the treatment of conditions in infants which are associated may be used in the treatment of conditions in infants which are associated with significant mortality or morbidity (such as pertussis or chlamydia), the benefit of erythromycin therapy needs to be weighed against the potential risk of developing IHPS. Parents and caregivers should be informed to contact their physician if vomiting and / or irritability with feeding occurs.
- In the event of severe acute hypersensitivity reactions, such as anaphylaxis, severe cutaneous adverse reactions (SCARs) [e.g. Stevens-Johnson Syndrome (SJS), toxic epidermal necrolysis (TEN), drug reaction with eosinophilia and systemic symptoms (DRESS) & acute generalised exanthematous pustulosis (AGEP)], Erogran should be discontinued immediately and appropriate treatment should be urgently initiated.

MAIN SIDE/ADVERSE EFFECTS

- Abdominal discomfort, diarrhoea, nausea, vomiting and ototoxicity (after high doses).
- Mild allergic reactions such as urticaria and other skin rashes.
- Serious allergic reactions including anaphylaxis.
- Hearing loss and/or tinnitus (patients with renal and hepatic insufficiency and on high dose).

Postmarketing Experience:

Gastrointestinal Disorders: infantile hypertrophic pyloric stenosis

Skin and Subcutaneous Tissue Disorders

Frequency not known: severe cutaneous adverse reactions (SCARs) including Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), drug reaction with eosinophilia and systemic symptoms (DRESS) & acute generalised exanthematous pustulosis (AGEP).

DRUG INTERACTION

- Concurrent use with aminophylline, oxtriphylline or theophylline may result in increased serum theophylline levels; dosage adjustments may be necessary.
- Co administration of lincomycins or penicillins is not recommended since erythromycins may interfere with their effects.

OVERDOSAGE

Symptoms of overdose

Nausea, vomiting, diarrhoea, prostration.

Treat overdosage by emesis or gastric lavage, if appropriate; with the necessary symptomatic and supportive measures, if required.

DOSAGE AND ADMINISTRATION

Adults:

- **Antibacterial**
Oral, the equivalent of erythromycin 400 mg every 6 hours or 800 mg every 12 hours.
- **Antibacterial (streptococcal prophylaxis)** - continuous prophylaxis of streptococcal infections in patients with a history of rheumatic heart disease and/or chorea.
Oral, the equivalent of erythromycin 400 mg every 12 hours
- **Antiprotozoal**
Oral, the equivalent of erythromycin 400 mg every 6 hours for ten to fourteen days.
- **Gonorrhoea (disseminated)**
Oral, the equivalent of erythromycin 800 mg every 6 hours for seven days.
- **Legionnaires' Disease**
Oral, the equivalent of erythromycin 400 mg to 1 gram every 6 hours.
- **Syphilis**
Oral, the equivalent of erythromycin 800 mg every 6 hours for 15 days (early syphilis) or for 30 days (late syphilis).

Usual adult prescribing limits:

Antibacterial the equivalent of erythromycin up to 4 grams daily.

Note: Doses up to the equivalent of 8 grams of erythromycin daily are apparently well tolerated.

Children:

- **Antibacterial**
oral, the equivalent of erythromycin 7.5 to 25 mg per kg of body weight every 6 hours
- **Antiprotozoal**
Oral, the equivalent of erythromycin 7.5 to 12.5 mg per kg of body weight every 6 hours for 10 to 14 days.

Note: 1.17 grams of erythromycin ethylsuccinate are equivalent to approximately 1 gram of erythromycin.

To reconstitute, add cooled boiled water until well dispersed, then add water to the appropriate mark.

Note: The information given here is limited. For further information, consult your doctor or pharmacist.

Storage: Store below 30°C. Protect from moisture. Refrigerate and use within 10 days after reconstitution.

Presentation/Packing:
Granules 200 mg/5 mL x 60 mL and 100 mL

Product Registration Holder: HOVID Bhd.
121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar),
30010 Ipoh, Perak, Malaysia.

Manufactured by: HOVID Bhd.
Lot 56442, 7 1/2 Miles, Jalan Ipoh / Chemor,
31200 Chemor, Perak, Malaysia.

Revision date: June 2025

