



○ SWKD
○ WKMS
○ WKMM
○ HU
○ HU (KK)

File Name : Medicinal Powder_Leaflet_Msia_02122020

Received : 02.12.2020

Amended : -

Output : -

Requestor : Shuying

PIC :

Designer : Nickles

Version : 00

Opened : 140w x 200h MM

Closed : -

Finished :

Colour : 1Cx1C





Three Legs Medicinal Powder Aspirin 600mg/sachet
Ubat Serbuk Cap Kaki Tiga Aspirin 600mg/bungkus
三脚标止痛退热粉阿司匹林600毫克/包

Three Legs Medicinal Powder Aspirin 600mg/sachet

Composition: Aspirin 600mg

Description: White powder, odourless or almost odourless with a slight acid taste.

Indication: For relief of headache, fever, toothache and minor pain.

Dosage:

Adult: 1 sachet every 4 hours. Maximum 6 sachets per day. To be taken after meal. Dissolve medicinal powder in a glass of water before taking.

Route of Administration: Oral

Contraindication: Patient with haemophilia. In patients where renal or hepatic functions is impaired.

Warning and Precaution:

Aspirin should be given with care to patient known to have a lesion of the gastric mucosa or patients with impaired renal or hepatic function. Daily use may potentiate oral anticoagulants and sulphonylurea hypoglycemic agents.

Not to be given to children under 16 years of age.

Drug Interaction:

The effects of the medicines or groups of substances below may be affected by concomitant treatment with Acetylsalicylic acid.

Enhanced effects ranging up to an increased risk of side effects:

Anticoagulants (e.g. coumarin, heparin) and thrombolytic drugs: acetylsalicylic acid can increase the risk of bleeding if taken before anticoagulant treatment.

Other platelet aggregation inhibitors (drugs that inhibit the clumping together of blood platelets), e.g. ticlopidine, clopidogrel: increased bleeding risk.

Drug product containing cortisone or cortisone- like substances (with the exception of products that are applied topically or cortisone replacement therapy for Addison's disease); increased risk of gastrointestinal side effects.

Alcohol: increased risk of gastrointestinal ulcers and bleeding.

Other analgesics and anti- inflammatory medicines (non- steroidal anti- inflammatory drugs) and other anti-rheumatic medicines in general: increased risk for gastrointestinal ulcers and bleeding.

Anti-diabetics: the blood glucose level can be reduces.

Digoxin (a drug to strengthen the heart muscle contraction).

Methotrexate (a drug used to treat cancer and certain rheumatic disorders).

Valproic acid (a drug used to treat convulsions) of the brain.

Selective Serotonin re- uptake inhibitors (certain medicines for treatment of depression) increased risk for gastrointestinal bleeding.

Certain antibiotics (sulphonamides and sulphonamide combinations, e.g. sulphamethoxazole/ trimethoprim).

Triiodothyronine (a medicine for treating hypothyroidism).

Weakening of effects:

Certain medicines that increase the excretion of urine (diuretics so-called aldosterone antagonists, e.g. spironolactone and canrenoate, and loop diuretics, e.g. furosemide).

Certain antihypertension drugs in particular angiotensin converting enzyme inhibitors).

Gout remedies that promote the excretion of uric acid (e.g. probenecid, benzbromarone).

Pregnancy and Lactation:

Pregnancy

Acetylsalicylic acid used during the third trimester of pregnancy should be avoided because of the known effects of NSAIDs on the fetal cardiovascular system. Salicylate have also been associated with alterations in maternal and neonatal hemostasis mechanisms, decreased birth weight, and with perinatal morality.

Lactation

Nursing mothers should avoid using Acetylsalicylic acid. Use of high doses may lead to rashes, platelet abnormalities, and bleeding in nursing infants.

Side Effects: Nausea, dyspepsia and vomiting. Irritation of the gastric mucosa and ulceration may occur. Aspirin increase the bleeding time and decrease platelet adhesiveness.

Symptoms and Treatment of Overdose:

Dizziness, tinnitus, deafness, sweating, nausea and vomiting, headache and mental confusion. In severe poisoning, hyperventilation, fever, restlessness, ketosis and respiratory alkalosis and metabolic acidosis results. Depression of the central nervous system may lead to coma, cardiovascular collapse and respiratory failure.

In acute salicylate overdose, the stomach should be emptied by inducing emesis or by aspiration and lavage. Patients with mild intoxication should be encouraged to drink plenty of fluids. In patients with more severe intoxication (plasma salicylate concentration above 500µg per ml in adults or 300µg per ml in children), forced alkaline diuresis may be required and should be continued until the plasma salicylate concentration is less than 350µg per ml in adults.

Pharmacodynamics:

Aspirin has analgesic and antipyretic as well as anti-inflammatory effects. It alleviates pain by inhibiting prostaglandin synthesis in the central nervous system and through a peripheral action by inhibiting the synthesis of prostaglandins that occurs in inflamed tissues to prevent the sensitization of the pain receptors to mechanical or chemical stimulation. Aspirin produces antipyrasis by acting centrally on the hypothalamic heat-regulating center by inhibiting prostaglandin synthesis in the hypothalamus to produce peripheral vasodilation, resulting in increased cutaneous blood flow, sweating and heat loss. Its anti-inflammatory activity is by inhibiting the synthesis of prostaglandin in the inflamed tissue.

Pharmacokinetics:

Absorption is rapid and complete following oral administration mostly from the upper small intestine. Food decrease the rate but not the extent of absorption. The plasma concentration reaches a peak in 1 to 2 hours, and the plasma half-life is about 2 to 3 hours. Aspirin is primarily metabolized in the liver. It is excreted in the urine as salicylic acid (10%) and as glucuronide conjugates (15%) and as salicyluric (75%) and gestisic acids (<1%). The rate of excretion of aspirin varies with the pH of the urine, increasing as the pH rises.

Storage Condition:

Store below 30°C. Protect from light and moisture. Keep out of reach of children.

Shelf Life: 3 years

Dosage Form: Powder

Packing Size: 0.95g x 12 Sachets

The information contained in this leaflet is limited. For further information, please consult your pharmacist or doctor.

Ubat Serbuk Cap Kaki Tiga Aspirin 600mg/bungkus

Komposisi: Aspirin 600mg

Deskripsi: Serbuk putih yang tidak berbau atau hampir tidak berbau dengan sedikit rasa berasid.

Indikasi: Untuk melegakan sakit kepala, demam, sakit gigi dan kesakitan yang ringan.

Cara Penggunaan:

Dewasa: 1 bungkus setiap 4 jam. Maksimum 6 bungkus sehari. Ambil selepas makan. Larutkan ubat serbuk ini di dalam segelas air dan minum.

Cara Pemberian Ubat: Oral

Kontraindikasi: Pesakit dengan haemophilia dan masalah hati atau buah pinggang.

Amaran dan Perhatian:

Aspirin perlu diberikan dengan berhati-hati untuk pesakit yang mempunyai iritasi mukosa gastrik atau pesakit yang mempunyai penjejasan pada buah pinggang atau hati. Penggunaan harian boleh meningkat kesan antikoagulan dan agen hipoglisemik sulfonulurea. Tidak boleh diberi kepada kanak-kanak berumur kurang daripada 16 tahun.

Interaksi dengan ubat lain: Sila rujuk kepada penerangan Bahasa Inggeris.

Kehamilan dan Penyusuan: Sila rujuk kepada penerangan Bahasa Inggeris.

Kesan Sampingan: Loya, dyspepsia dan muntah mungkin berlaku. Iritasi pada mukosa gastrik dan ulser boleh berlaku. Aspirin meningkatkan pendarahan dan menurunkan kepekatan platelet.

Symptom dan Rawatan untuk Dos Berlebihan: Sila rujuk kepada penerangan Bahasa Inggeris.

Farmakodinamik: Sila rujuk kepada penerangan Bahasa Inggeris.

Farmakokinetik: Sila rujuk kepada penerangan Bahasa Inggeris.

Keadaan Penyimpanan:

Simpan di bawah 30°C. Lindungi dari cahaya dan kelembapan. Jauhkan daripada capaian kanak-kanak.

Tempoh Hayat: 3 tahun

Bentuk Dos: Serbuk

Saiz Bungkus: 0.95g x 12 Bungkus

Maklumat yang terkandung di dalam risalah ini adalah terhad. Untuk maklumat yang terperinci, sila hubungi ahli farmasi atau doktor.

三脚标止痛退热粉阿司匹林600毫克/包

成份: 阿司匹林600毫克

外观: 带有微酸味，无臭或近乎无臭白色粉末。

主治: 能有效地解除头痛、发烧、牙痛及轻微疼痛。

用量:

成人：每4小时1包。每日最多可服用6包。饭后服用。将药粉溶解在1杯开水后服食。

给药途径：口服。

禁忌: 血友病患者，肾功能或肝功能衰竭病患者。

注意及防范：

阿司匹林应谨慎给予胃粘膜病变或肾功能或肝功能受损的患者。日常使用中，可能会增加口服抗凝血药物和磺胺类降糖药。16岁以下儿童不宜服用。

药物相互作用：请参阅英文版本。

怀孕和哺乳：请参阅英文版本。

副作用: 头晕、耳鸣、耳聋、出汗、恶心、呕吐、头痛和 精神混乱。当严重中毒时，会导致过度换气、发热、烦躁不安、酮症和呼吸性碱中毒和代谢性酸中毒。中枢神经系统被压制，可能导致昏迷，心血管功能衰竭和呼吸衰竭。

药物过量症状及治疗：请参阅英文版本。

药效学：请参阅英文版本。

药代动力学：请参阅英文版本。

贮存：储藏低于30°C。避免阳光直射及潮湿处。远离儿童。

有效期：3年

剂型：粉末

包装：0.95克 x 12包

此说明书仅属略述，欲知更详细资料，可咨询您的药剂师或医生。

Manufactured by Product Registration Holder:
Syarikat Wen Ken Drug Sdn. Bhd.
No. 24, Jalan Lambak, Taman Johor,
81200 Johor Bahru, Johor, Malaysia.

Pengedar Singapura/ Singapore Distributor:
Wen Ken Marketing (S) Pte Ltd
2, Alexandra Road, #02-08, Delta House,
Singapore 159919.

Date of Revision: 02-12-2020

...