

**ATEBLOC TABLET 50mg**

Yellow, round, biconvex, top-scored tablet

ATEBLOC TABLET 100mg

Orange, round, flat, top-scored tablet

Content

Atebloc Tablet 50mg : Atenolol B.P. 50mg/tablet

Atebloc Tablet 100mg : Atenolol B.P. 100mg/tablet

Pharmacology

Atenolol, a cardioselective beta-1-adrenergic blocker, has antihypertensive, antianginal, and antiarrhythmic properties. Its antianginal effect is probably due to reduced myocardial oxygen requirements, antiarrhythmic effect due to blockage of adrenergic stimulation of cardiac pacemaker potentials; and antihypertensive effect may include reduced cardiac output, decreased sympathetic outflow to peripheral vasculature, and inhibition of renin release by kidney.

Oral absorption is complete, hepatic metabolism is minimal, and elimination is by renal excretion.

Indications

For treatment of hypertension, angina pectoris and cardiac arrhythmias, and prophylactic treatment of myocardial reinfarction.

Adverse Effects

Fatigue, depression, cold extremities, skin rash (allergic reaction), diarrhoea, dizziness, headache, nausea or vomiting, and weakness may occur occasionally.

Precautions

- Caution in patients with history of allergy, mental depression, bronchial asthma, congestive heart failure, hypoglycaemia, or renal function impairment.
- Check with physician before discontinuing medication; gradual dosage reduction may be necessary in the treatment of coronary artery disease.
- Safety for use in pregnancy and lactation has not been established.

Contraindications

Avoid in patients with overt cardiac failure, cardiogenic shock, 2nd-or-3rd-degree atrioventricular (AV) block, or sinus bradycardia (heart rates less than 45 beats per minute).

Drug Interactions

- Concurrent use with general anaesthetics may increase the risk of myocardial depression and hypotension.
- Co-administration of calcium channel blocking agents or clonidine may potentiate antihypertensive effects.
- Concomitant therapy with sympathomimetic amines having beta-adrenergic stimulant activity may result in mutual inhibition of the therapeutic effects.

Overdosage

Signs of overdose include slow heartbeat, severe dizziness, fainting, difficulty in breathing, bluish-coloured fingernails or palms of hands, or seizures.

For treatment of overdose, in addition to gastric lavage, supportive and symptomatic therapy are required to treat bradycardia, premature ventricular contraction, cardiac failure, hypotension, seizures, and bronchospasm.

Dosage and Administration

Adults - Antianginal : Oral, initially 50mg once a day, the dosage being increased gradually to 100mg after one week if necessary and tolerated. Some patients may require up to 200mg per day.

- Antihypertensive : Oral, initially 25 to 50mg once a day, the dosage being increased to 50 to 100mg a day after two weeks if necessary and tolerated.

Note : If creatinine clearance is 15 to 35 mL/min/1.73m², the maximum dose is 50mg/day; if it is less than 15mL/min/1.73m², it is 50mg every second day.

Children - Dosage has not been established.

Storage : Keep container tightly closed. Store in a cool, dry place, below 30°C. Protect from light.

Presentation/ Packing : Tablet 50mg x 1000's
Tablet 100mg x 1000's

Shelf-life : 3 years

Manufactured & : Prime Pharmaceutical Sdn. Bhd.

Distributed by : 1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah,
14000 Bukit Mertajam, Penang, Malaysia.