



MYOFLEX TABLET

Content: Each tablet contains: Orphenadrine citrate BP/USP 35mg, Paracetamol BP/USP 450mg

Pharmacology

Orphenadrine reduces voluntary muscle spasm by a central effect it has analgesic activity, which may contribute to its skeletal muscle relaxant properties. Paracetamol, an analgesic and antipyretic, acts by inhibiting prostaglandin synthesis in the central nervous system and to a lesser extent, through a peripheral action by blocking pain impulse generation. It is readily absorbed orally, metabolised in the liver, and excreted in the urine.

Indications:

For relief of muscle spasm associated with acute painful musculoskeletal conditions.

Adverse Effects / Undesirable Effect:

Blurred vision, dry mouth, nausea or vomiting, vertigo, skin rash or reactions, headache, dizziness, drowsiness, constipation, increased intraocular pressure, confusion, excitement, difficult urination and gastrointestinal pains may occur occasionally. Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome/Toxic Epidermal Necrolysis have been reported.

Warning:

Allergy alert : Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Precautions

- Caution in patients with cardiac disease or arrhythmias or tachycardia, CNS depression, hepatic or renal function impairment , and urinary retention.
- Caution when driving or doing jobs requiring alertness because of possible drowsiness.
- Safety for use in pregnancy and lactation has not been established.

Contraindications

Avoid in patients with achalasia, bladder neck obstruction, glaucoma, myasthenia gravis, prostatic hypertrophy, or pyloric or duodenal obstruction.

Drug Interactions

- Concurrent use with CNS depression-producing medications may result in additive CNS depressant effects; dosage of one or both agents should be reduced.
- Co-administration of antimuscarinics or other medications with antimuscarinic action may intensify antimuscarinic effects.
- Risk of hepatotoxicity with prolonged use of high doses of this medication may be increased in alcoholics or in patients regularly taking other hepatotoxic medications or hepatic enzyme inducers.

Overdosage

The safety of continuous long-term administration of orphenadrine has not been established.

Signs of paracetamol overdosage include diarrhoea, increased sweating, anorexia, nausea or vomiting, stomach or abdominal cramps, hepatic encephalopathy, coma, metabolic acidosis, renal tubular necrosis, cardiac arrhythmias and cardiovascular collapse.

Treatment consists of emesis or gastric lavage; instituting supportive treatment, including maintaining fluid and electrolyte balance; administering acetylcysteine as soon as possible or within 10 to 12 hours after ingestion of the overdose; and instituting hemodialysis or hemoperfusion if too much time has elapsed.

Dosage and Administration

- | | |
|------------------------|---|
| Adults | : Oral, 1 to 2 tablets three times a day, adjusted according to the patient's response, |
| Children | : Dosage has not been established. |
| Storage | : Keep container tightly closed. Store in a cool, dry place below 30°C. Protect from light. |
| Presentation / Packing | : Tablet X 1000's |
| Shelf life | : 3 years |
| Manufactured & | : PRIME PHARMACEUTICAL SDN. BHD. |
| Distributed by | : 1505, Lrg Perusahaan Utama 1, Tmn Perindustrian Bkt Tengah, 14000 Bkt Mertajam, Pg, Malaysia. |