

VOREN® GEL

Voren, the brand of Diclofenac, is an anti-inflammatory and analgesic agent.

Ingredient(s):

Each gram contains:

Diclofenac Acid Diethylamine Salt 11.6mg
(eq. to 10mg of Diclofenac Sodium)

Pharmacology (Summary of Pharmacokinetics and Pharmacodynamics):

Pharmacodynamics

- a. Voren Gel is an anti-inflammatory agent and analgesic preparation designed for external application. It contains a quantity of active substances equivalent to 1% Diclofenac Sodium. Voren Gel can be easily rubbed onto the skin. Due to its aqueous-alcoholic base it exerts a soothing and cooling effect. Thus it is effective in the relief of pain and swelling due to rheumatosis of inflammation.
- b. Voren has major anti-inflammatory properties, which apparently are related to the inhibition of cyclo-oxygenase, a decrease in prostaglandin synthesis, and an increase in the reuptake of Arachidonic Acid into cellular pools.
- c. Voren Gel is suitable for use in inflammatory and degenerative rheumatic diseases, as well as for the treatment of post-traumatic inflammation and swelling.
Diclofenac Sodium is a non-steroidal-type anti-inflammatory agent and is widely used clinically because of its strong analgesic, antipyretic and anti-inflammatory effects.

Pharmacokinetics

- a. Voren Gel to the skin in healthy subjects as compared with oral administration of Voren coated tablet is equivalent to approximately 6% of the dose applied.
- b. Diclofenac Sodium is absorbed from the gastrointestinal tract. Peak plasma concentrations occur about 2 hours after ingestion of enteric coated tablets. At therapeutic concentrations it is more than 99% bound to plasma protein. Diclofenac Sodium undergoes extensive biotransformation in the liver during the first passage through which about 50% is metabolized. Approximately 1/3 of the dose is excreted in the bile and approximately 2/3 via the kidneys in the form of glucuronides. Then mean terminal elimination half life of Diclofenac Sodium is 1.2 to 1.8 hours and metabolite possesses about 1/30 of the anti-inflammatory activity of parent drug (Diclofenac Sodium).

Indication(s):

Topical application in the treatment of inflammatory conditions or pain caused by chronic rheumatoid arthritis, degenerative arthritis and soft tissue arthritis.

Dosage and Administration:

Depending on the size of the painful site to be treated, apply 2 – 4g of Voren Gel 3 to 4 times daily to the affected areas and rub in gently. After assessing the risk/ benefit ratio in each individual patient, the lowest effective dose for the shortest possible duration should be used.

Mode Of Administration:

Listed in recommended dosage.

Contraindication(s):

It is contraindicated in patients with a history of hypersensitivity to Diclofenac, Acetylsalicylic acid and other non-steroidal anti-inflammatory drug.

Precaution(s) / Warning(s):

1. It should be applied only to intact skin surfaces, and not to skin wounds or open injuries.
2. It should not be allowed to come into contact with the eyes or with mucous membranes.
3. Since no experience has been acquired with Voren Gel in pregnant women, it is not recommended for use during pregnancy.
4. NSAIDs may very rarely cause serious cutaneous adverse events such as exfoliative dermatitis, toxic epidermal necrolysis (TEN) and Stevens-Johnson Syndrome (SJS), which can be fatal and without warning. These serious adverse events are idiosyncratic and are independent of dose or duration of use. Patients should be advised of the signs and symptoms of serious skin reaction and to consult their doctor at the first appearance of a skin rash or any other sign of hypersensitivity.
5. When Voren Gel is applied to relatively large areas of skin and over a prolonged period, the possibility of systemic side effects cannot be completely excluded. In such cases, the product information on Voren should be consulted.
6. Observational studies have indicated that non-selective NSAIDs may be associated with an increased risk of serious cardiovascular events, principally myocardial infarction, which may increase with dose or duration of use. Patients with cardiovascular disease or cardiovascular risk of an adverse cardiovascular event in patient taking NSAID, especially in those with cardiovascular risk factors, the lowest effective dose should be used for the shortest possible duration. There is no consistent evidence that the concurrent use of aspirin mitigates the possible increased risk of serious cardiovascular thrombotic events associated with NSAID use.
7. NSAIDs may lead to the onset of new hypertension or worsening the pre-existing hypertension and patients taking anti-hypertensive with NSAIDs may have an impaired anti-hypertensive response. Caution is advised when prescribing NSAIDs to patients with hypertension. Blood pressure should be monitored closely during initiation of NSAID treatment and at regular intervals thereafter.
8. Fluid retention and oedema have been observed in some patients taking NSAIDs, therefore caution is advised in patients with fluid retention or heart failure.
9. All NSAIDs can cause gastrointestinal discomfort and rarely serious, potentially fatal gastrointestinal effects such as ulcers, bleeding and perforation which may increase with dose or duration of use, but can occur at any time without warning. Caution is advised in patients with risk factors for gastrointestinal events e.g. the elderly, those with a history of serious gastrointestinal events, smoking and alcoholism. When gastrointestinal bleeding or ulceration occurs in patients receiving NSAIDs, the drug should be withdrawn immediately. Doctors should warn patients about signs and symptoms of serious gastrointestinal toxicity. The concurrent use of aspirin and NSAIDs also increases the risk of serious gastrointestinal adverse events.

Interactions With Other Medicaments:

None has been mentioned

Pregnancy and Lactation:

Since no experience has been acquired with Voren Gel in pregnant women, it is not recommended for use during pregnancy.

Side Effect(s) / Adverse Reaction(s):

Voren Gel is usually well tolerated. Itching, reddening, or smarting of the skin or outbreak of a skin rash may occasionally occur.
Dermatological : Occasional rashes or skin eruptions.
Cases of hair loss, bullous eruptions, erythema multiforme, Steven's-Johnson Syndrome, toxic epidermal necrolysis (Lyell's syndrome) and photosensitivity reactions have been reported.

Symptoms and Treatment for Overdosage, and Antidote(s):

In the event of significant systemic side effects occurring as a result of improper use or an accidental overdosage (eg. in children), general therapeutic measures of the kind normally adopted in order to treat poisoning with non-steroidal anti-inflammatory drug should be instituted. Hemodialysis and peritoneal dialysis may be of little value.

Shelf-Life:

3 years from the date of manufacture.

Storage Condition(s):

Keep in an airtight container. Store at temperature below 30°C. Protect from light and moisture.

Product Description and Packing(s):

A colorless to white color gel
Aluminium tube of 20g.

WARNINGS

RISK OF GI ULCERATION, BLEEDING AND PERFORATION WITH NSAID

Serious GI toxicity such as bleeding, ulceration and perforation can occur at any time, with or without warning symptoms, in patients treated with NSAID therapy. Although minor upper GI problems (eg. dyspepsia) are common, usually developing early in therapy, prescribers should remain alert for ulceration and bleeding in patients treated with NSAIDs even in the absence of previous GI tract symptoms.

Studies to date have not identified any subset of patients not at risk of developing peptic ulceration and bleeding. Patients with prior history of serious adverse events and other risk factors associated with peptic ulcer disease (eg. alcoholism, smoking, and corticosteroid therapy) are at increased risk. Elderly or debilitated patients seem to tolerate ulceration or bleeding less than other individuals and account for most spontaneous reports for fatal GI events.

VOREN® GEL

Bahan aktif:

Setiap gram mengandungi:
Diclofenac Acid Diethylamine Salt 11.6mg
(setara dengan 10mg Diclofenac Sodium)

Indikasi:

Rawatan topical dalam keadaan inflamasi atau kesakitan akibat arthritis.

Dos:

Bergantung pada keluasan yang terlibat, sapu 2 - 4g Voren gel 3 - 4 kali sehari.

非 炎 乳 膠

成份:

每公克中含有:
Diclofenac Acid Diethylamine Salt 11.6mg
(相等於 Diclofenac Sodium 10mg)

適應症:

類風濕性關節炎、關節周圍疾患及各種炎症之消炎鎮痛局部治療。

用法、用量:

每天3~4次徐徐塗擦2~4公克本劑於皮膚患處;視發炎範圍斟酌塗擦之。



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679