

PRODUCT LITERATURE

PANAMOL SUSPENSION 250MG/5ML

Each 5ml contains:

Paracetamol 250 mg

Preservatives:

Methyl Paraben 0.1% w/v
Propyl Paraben 0.01% w/v

Product description:

Orange coloured viscous suspension with sweet taste and fruity odour.

Pharmacodynamics

Paracetamol is an antipyretic analgesic. The mechanism of action is probably similar to that of aspirin and dependent on the inhibition of prostaglandin synthesis. This inhibition appears, however to be on a selective basis.

Pharmacokinetics

Paracetamol is rapidly and almost completely absorbed from the gastrointestinal tract. The concentration in plasma reaches a peak in 30 to 60 minutes and the half-life in plasma is 1 to 4 hours after therapeutic doses. Paracetamol is relatively uniformly distributed throughout most body fluids. Binding of the drug to plasma proteins is variable; 20 to 50% may be bound at the concentrations encountered during acute intoxication. Following therapeutic doses 90 to 100% of the drug may be recovered in the urine within the first day. However, practically no Paracetamol is excreted unchanged, and the bulk is excreted after hepatic conjugation.

Indication

Analgesic and anti-pyretic.

Recommended Dose

To be given 3 or 4 times a day.
Children (6-12 years): 5ml – 10 ml (1-2 tsp)
(1-5 years): 2.5ml – 5 ml (1/2– 1 tsp)

Route of Administration

Oral

Contraindications

Hypersensitivity to Paracetamol or any of the other constituents.

Warnings and Precautions

This preparation contains PARACETAMOL.
Do not take any other Paracetamol containing medicines at the same time.

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Care is advised in the administration of Paracetamol to patients with severe renal or severe hepatic impairment. The hazards of overdose are greater in those with (non-cirrhotic) alcoholic liver disease.

- Do not overfill the spoon.
- Always use the spoon supplied with the pack.
- Do not give more than 4 doses in any 24 hour period.
- Leave at least 4 hours between doses.
- Do not give this medicine to your child for more than 3 days without speaking to your doctor or pharmacist.
- As with all medicines, if your child is currently taking any medicine consult your doctor or pharmacist before taking this product.

Interactions with Other Medicaments

The speed of absorption of Paracetamol may be increased by metoclopramide or domperidone and absorption reduced by cholestyramine. The anti-coagulant effect of warfarin and other coumarins may be enhanced by prolonged regular use of Paracetamol with increased risk of bleeding; occasional doses have no significant effect.

Drugs which induce hepatic microsomal enzymes such as alcohol. Concomitant barbiturates and tricyclic antidepressants may increase the hepatotoxicity of Paracetamol particularly after overdose.

Anti-convulsion or oral steroid contraceptives have the ability to reduce serum levels of Paracetamol by liver enzyme induction.

Pregnancy and Lactation

Epidemiological studies in human pregnancy have shown no ill effects due to Paracetamol used in the recommended dosage, but patients should follow the advice of their doctor regarding its use. A large amount of data on pregnant women indicates neither malformative, nor feto/neonatal toxicity. Epidemiological studies on neurodevelopment in children exposed to Paracetamol in utero show inconclusive results. If clinically needed, Paracetamol can be used during pregnancy however it should be used at the lowest effective dose for the shortest possible time and at the lowest possible frequency.

Paracetamol is excreted in breast milk but not in a clinically significant amount. Available published data does not contraindicate breast-feeding.

Side Effects

Very rare cases of serious skin reactions have been reported. Adverse effects of Paracetamol are rare but hypersensitivity including skin rash may occur. There have been reports of blood dyscrasias including thrombocytopenia and agranulocytosis, but these

were not necessarily causally related to Paracetamol. With prolonged use or overdosage, hepatic necrosis, acute pancreatitis and nephrotoxicity have been reported.

Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome/Toxic Epidermal Necrolysis have been reported.

Symptoms and Treatment of overdose

Immediate treatment is essential in the management of Paracetamol overdose. Despite a lack of significant early symptoms, patients should be referred to hospital urgently for immediate medical attention and any patient who had ingested around 7.5g or more of Paracetamol in the preceding 4 hours should undergo gastric lavage. Administration of oral methionine or intravenous N-acetylcysteine which may have a beneficial effect up to at least 48 hours after the overdose may be required. General supportive measures must be available.

Symptoms of Paracetamol overdosage in the first 24 hours are pallor, nausea, vomiting, anorexia and abdominal pain. Liver damage may become apparent 12 to 48 hours after ingestion. Abnormalities of glucose metabolism and metabolic acidosis may occur. In severe poisoning, hepatic failure may progress to encephalopathy, coma and death. Acute renal failure with acute tubular necrosis may develop even in the absence of severe liver damage. Cardiac arrhythmias and pancreatitis have been reported.

Liver damage is possible in adults who have taken 10g or more of Paracetamol.

Packing

Packed in plastic bottle of 60 ml and 90 ml.

Storage Condition

Store below 30°C in a dry place, protected from direct light.
Keep away from reach of children.
SHAKE WELL BEFORE USE

Expiry Date

2 years from date of manufacture

Name and Address of Product Registration Holder/ Distributor:

ZONTRON PHARMACEUTICALS SDN.BHD. (445695-T)
Lot 10 & 11, PERDA Industrial Park,
Lorong IKS Simpang Ampat B,
14100 Simpang Ampat, S.P.S.,
Pulau Pinang, Malaysia

Name and address of Manufacturer:

TERAPUTICS SDN. BHD. (590500-W)
Lot 10 & 11, PERDA Industrial Park,
Lorong IKS Simpang Ampat B,
14100 Simpang Ampat, S.P.S.,
Pulau Pinang, Malaysia

Malaysia Drug Registration no:

MAL19940171XCZ

Date of Revision: 26/08/2025