

PRODUCT LITERATURE

PANAMOL 500MG TABLET

Each tablet contains:

Paracetamol 500 mg

Preservative:

Sodium Benzoate 0.1% w/w

Product description:

Round, pink coloured tablet, with the word "ZONTRON/500" embossed on one side.

Pharmacodynamics

Paracetamol is a centrally acting analgesic and antipyretic with minimal anti-inflammatory properties.

Analgesic

The mechanism of analgesic action has not been fully determined. Paracetamol may act predominantly by inhibiting prostaglandin synthesis in the central nervous system (specifically cyclooxygenase (COX)-2) and, to a lesser extent, through a peripheral action by blocking pain-impulse generation.

The peripheral action may also be due to inhibition of prostaglandin synthesis or to inhibition of the synthesis or actions of other substances that sensitize pain receptors to mechanical or chemical stimulation.

Antipyretic

Paracetamol act centrally on the hypothalamic heat-regulating center to produce peripheral vasodilatation resulting in increased blood flow through the skin, sweating and heat loss. Paracetamol reduces fever by inhibiting the formulation and release of prostaglandins in the CNS and by inhibiting endogenous pyrogens at the hypothalamic thermo regulator center.

Pharmacokinetics

Following oral administration Paracetamol is rapidly absorbed.

Paracetamol absorption takes place mainly in the small intestine and therefore the rate of absorption is depending on the rate of gastric emptying. It has been shown that drugs which delay gastric emptying also delay the absorption of Paracetamol whereas metoclopramide (a drug which increases the rate of gastric emptying) accelerates absorption of the analgesic through the total amount absorbed doses not increase.

The presence of food in the stomach has also been reported to reduce the rate of absorption of Paracetamol. Alterations in gastric pH have no appreciable effect on Paracetamol absorption. During absorption, the amount of Paracetamol does not affect gastric mucosal permeability and does not produce mucosal bleeding. Peak plasma concentrations are reached 1 hour after absorption. The plasma half-life is 1 to 3 hours.

Paracetamol penetrates the brain and is present in breast milk of human.

Paracetamol is metabolized by the microsomal enzyme system of the liver. This metabolism is mainly to the glucuronide and sulphate conjugates, accounting for approximately 49% and 26% of the ingested dose respectively. About 4% is excreted as free Paracetamol. Other minor pathways include the production of catechol derivatives and cysteine conjugates (via glutathione). Paracetamol excretion is rapid and occurs via the urine.

Indication

- For the relief of fever
- For the relief from mild to moderate including: headache, migraine, backache, musculoskeletal pain, myalgia and neuralgia, dysmenorrhea, pain of osteoarthritis, toothache, pain after dental procedures / tooth extraction, pain after vaccination and the discomfort from colds, influenza and sore throats.

Recommended Dose

Adults & Children Aged 12 Years and Over

500mg to 1g paracetamol, taken every 4-6 hours as required up to a maximum of 4g daily

Children

6 To 11 Years: 250mg - 500mg every 4 -6 hours as required

Maximum daily dose: 60mg / kg presented in divided doses of 10 - 15 mg / kg throughout 24 hour period

Route of Administration

Oral

Contraindications

Hypersensitivity to Paracetamol or any of the other ingredients / components of the product. Severe and active hepatic impairment.

Warnings and Precautions

This preparation contains PARACETAMOL.
Do not take any other Paracetamol containing medicines at the same time.

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

- Keep out of reach of children
- Do not take if allergic to Paracetamol
- Patients should contact their health care provider if symptoms persist (if the pain lasts for more than 10 days, if there is redness or fever lasts more than 3 days)
- Paracetamol should be given with care to patients with impaired kidney or liver function
- Larger doses should be avoided in patients with hepatic impairment. Paracetamol overdose may harm the liver.
- Do not exceed recommended dose.
- It should be given with care to patients with alcohol dependence.
- Paracetamol provides symptomatic relief only, additional therapy to treat the cause of the pain or fever should be instituted when necessary.

Effects on Ability to Drive and Use Machines

It is unlikely to impair a patient’s ability to driver or use machinery

Interactions with Other Medicaments

The anticoagulant effect of warfarin and other coumarins may be enhanced by prolonged regular daily use of Paracetamol with increased risk of bleeding; occasional doses have no significant effect.

Pregnancy and Lactation

Use in Pregnancy

- Considered to be the analgesic of choice in pregnant patients
- Although it crosses placenta, Paracetamol is considered to be safe in normal therapeutic doses for short term use as a minor analgesic / antipyretic in pregnancy.

Use in Lactation

- Excreted in breast milk
- Maternal ingestion of Paracetamol in normal therapeutic doses does not appear to present a risk to the nursing infant.

Side Effects

Adverse effects of Paracetamol are rare and usually mild, although haematological reactions have been reported. Skin rashes and other hypersensitivity reactions occur occasionally.

Symptoms and Treatment of overdose

Symptoms

Toxic symptoms include vomiting, abdominal pain, hypotension and sweating. The most serious adverse effects of acute overdose of Paracetamol is a dose-dependent, potentially fatal hepatic necrosis. Clinical and laboratory evidence of hepatotoxicity may be delayed for up to one week. Major manifestations of liver failure such as jaundice, hypoglycemia and metabolic acidosis may take at least 3 days to develop.

Treatment

In cases of overdose, methods of reducing the absorption of ingested drug are important. Gastric lavage is essential even if several hours have elapsed. Prompt administration of 50g activated charcoal and 500mL iced mannitol 20% by mouth, may reduce absorption. If the history suggests that 15g Paracetamol or more has been ingested, administer one of the following antidotes:

Acetylcysteine 20% i.v.: Administer intravenously, 20% acetylcysteine (Parvolex, Glaxo) immediately without waiting for positive urine test or plasma level results : initial dose of 150mg / kg over 15 minutes, followed by continuous infusion of 50mg / kg in 500mL 5% glucose / dextrose over 4 hours and 100mg / kg in 1L 5% glucose / dextrose over 16 hours;

OR

Oral Methionine: 2.5g immediately followed by three further doses of 2.5g at four hourly intervals. For a 3 year old child, 1g methionine every four hours for four doses has been used;

OR

Oral Acetylcysteine 5%: 140mg / kg as a loading dose, then 70mg / kg every 4 hours for a total of 17 maintenance doses. If more than ten hours have elapsed since the overdosage was taken, the antidote may be ineffective.

Packing

Blister pack of 3x10’s, 10x10’s, 50x10’s and 100x10’s.

Storage Condition

Store below 30°C in a dry place, protected from direct light.

Keep away from reach of children.

Expiry Date

3 years from date of manufacture

Name and Address of Product Registration Holder/ Distributor:

ZONTRON PHARMACEUTICALS SDN.BHD. (445695-T)

Lot 10 & 11, PERDA Industrial Park,

Lorong IKS Simpang Ampat B,

14100 Simpang Ampat, S.P.S.,

Pulau Pinang, Malaysia

Name and address of Manufacturer:

TERAPUTICS SDN. BHD. (590500-W)

Lot 10 & 11, PERDA Industrial Park,

Lorong IKS Simpang Ampat B,

14100 Simpang Ampat, S.P.S.,

Pulau Pinang, Malaysia

Malaysia Drug Registration no:

MAL19940208XCZ

Date of Revision: 29/10/2025