

98 mm

98 mm

Melegakan Sakit Kepala

Uphamol 650



Uphamol 650 mengandungi acetaminophen, satu bahan aktif yang berkesan sebagai analgesik dan antipiretik. Ia boleh diambil untuk melegakan simptom sakit-sakit kepala termasuk migrain dan sakit kepala akibat ketegangan.

Uphamol 650 juga boleh diambil untuk melegakan kesakitan akibat selsema dan demam selsema.

DESKRIPSI

Pil berdiameter 13.5mm, berbentuk bulat, rata, bertanda 'mahkota dengan UPHAMOL' dan '650' berwarna biru muda.

Setiap tablet mengandungi acetaminophen 650mg.

SUKATAN

Dewasa dan kanak-kanak 12 tahun ke atas : 1 tablet, setiap 4 hingga 6 jam, (jangan melebihi 6 tablet sehari) Jika sakit berlarutan, sila jumpa doktor.

Untuk rawatan jangka panjang, jangan melebihi 4 tablet dalam jangkamasa 24 jam.

KEADAAN PENYIMPANAN

Jangan gunakan Uphamol 650 yang sudah tamat tempoh penggunaannya. Sila pastikan ubat yang telah dibuang dijauhkan daripada kanak-kanak.

KESAN SAMPINGAN

Bila digunakan mengikut arahan, kesan sampingan daripada Uphamol 650 jarang berlaku. Radang kulit dan lain-lain reaksi hipersensitif telah dilaporkan sekali-sekala.

KONTRAINDIKASI

Uphamol 650 tidak boleh digunakan jika anda alah kepada acetaminophen.

AMARAN

Jika sakit berlarutan, sila jumpa doktor atau ahli farmasi anda.

Ubat ini mengandungi PARACETAMOL. Pastikan anda tidak makan ubat demam, sakit kepala atau ubat-ubatan lain yang juga mengandungi PARACETAMOL pada masa yang sama.

BUNGKUSAN

Dalam pek blister 20's, 180's, 500's dan pek sachet 1's dan 30 x 1's.

Simpan di tempat yang kering di bawah 30°C.
Jauhi daripada kanak-kanak.
Jangka hayat : Sila rujuk kotak.

108 mm

专治头痛

Uphamol 650



Uphamol 650 含有 acetaminophen 成份, 它以有效的止痛及退热功能见称. 它主治各种头痛病症, 其中包括偏头痛和压力头痛。

Uphamol 650 也可作为主治头痛、疼痛、伤风、感冒及发烧等的良药。

成份:

13.5毫米直径, 圆, 扁注明“冠形与UPHAMOL”和“650”淡蓝色药片
每片含有650mg acetaminophen.

服法:

成人或十二岁以上的儿童 : 每4-6小时服用一片, (每日不可服用超过六片)。如病痛持续数天, 敬请看医生。

长期服用者在24小时内不可服用超过4片。

贮存:

不要保留过期的Uphamol 650药片, 并确保丢弃的药物远离儿童。

副作用:

Uphamol 650 属温和药物, 它所可能引起的副作用可谓微之又微。服用 acetaminophen 可能引起轻微皮肤瘙痒及其他敏感反应。

禁忌:

对 acetaminophen 过敏者不可服用 Uphamol 650。

警告/留意:

如果疼痛持续, 请即询问您的医生。

此药片含有PARACETAMOL成份。
不可同时服用任何其他含PARACETAMOL成份的药物。

包装:

备有20's, 180's 及 500's 片装和1's 及 30 x 1's 包装

收藏于乾燥, 摄氏30°C以下之处。
置于儿童不能触及之处。
有效服用期限: 请参阅盒子。

TREATMENT OF OVERDOSAGE

First empty stomach via induction of emesis or gastric lavage. It is recommended that acetylcysteine administration be instituted as soon as possible after ingestion of an overdose has been reported without waiting for the results of plasma acetaminophen concentration or other laboratory tests. Acetylcysteine is most effective if treatment is started within 10 to 12 hours after ingestion of the overdose; however, it may be of some benefit if treatment is started within 24 hours.

For oral administration, the recommended adult dose of acetylcysteine is 140mg per kg of body weight every 4 hours for 17 doses. Each dose should be diluted to a 5% solution with cola, grape fruit juice, orange juice or water prior to administration because of its unpleasant odor and its irritating and sclerosing properties. Any dose vomited within 1 hour of administration must be repeated. If necessary, the antidote may be given via duodenal intubation. Plasma acetaminophen concentration should be determined and liver function test should be performed to ascertain the status of the patient. Renal and cardiac function should be monitored and appropriate therapy instituted if necessary.

Supportive treatment consists of maintaining fluid and electrolyte balance, correcting hypoglycemia and administering vitamin K1 (if prothrombin time ratio exceeds 1.5) and fresh frozen plasma or clotting factor concentrate (if prothrombin ratio exceeds 3.0).

PHARMACODYNAMIC

Paracetamol has analgesic and antipyretic activities. The analgesic activity is brought about by the inhibition of the synthesis of prostaglandins. As for the antipyretic activity, paracetamol act centrally on the hypothalamic heat-regulating centre to produce peripheral vasodilation resulting in increased cutaneous blood flow, sweating and heat loss.

PHARMACOKINETICS

Paracetamol is rapidly and almost completely absorbed after oral administration. It is distributed into most body tissues. It crosses the placenta and is present in breast milk. Approximately 90-95% of a dose is metabolised in the liver, primary by conjugation with glucuronic acid, sulphur acid, and cysteine. Peak concentration can be achieved within 0.5-2 hours. Paracetamol is excreted primarily as conjugates via the kidney.

PACKING

In blister pack of 20's, 180's and 500's. Also available in sachet pack of 1's and 30 x 1's.

Store in a dry place below 30°C.
Keep out of reach of children.
Shelf life : Please refer to box.

STORAGE

Do not keep expired Uphamol 650 tablets. Be sure that any discarded medicine is out of reach of children.

Product Registration Holder :

DUOPHARMA MANUFACTURING (BANGI) SDN. BHD.
Lot No. 2, 4, 6, 8 & 10, Jalan P/7, Section 13, Bangi Industrial Estate, 43650 Bandar Baru Bangi, Selangor, Malaysia.



Manufacturer:

DUOPHARMA MANUFACTURING (BANGI) SDN. BHD.
Lot No. 2 & 4, Jalan P/7, Section 13, Bangi Industrial Estate, 43650 Bandar Baru Bangi, Selangor, Malaysia.

1500008811 N.1

Revision Date: July 2021

Relieves Headaches

Uphamol 650



Uphamol 650 contains acetaminophen which is a known effective analgesic and antipyretic. It can be used to relieve the symptoms of various types of headaches including migraine and tension-type headache. Uphamol 650 can also be used to relieve pain due to colds and flu and to reduce fever.

DESCRIPTION

A 13.5mm diameter, round, flat marked 'crown-shaped with UPHAMOL' and '650' light blue tablet. Each tablet contains acetaminophen 650mg.

Route of administration: Oral.

DOSAGE

Adult and children above 12 years: 1 tablet every 4 to 6 hours (not more than 6 tablets in one day). If symptoms persist, consult the doctor.

For long term treatment, not more than 4 tablets should be taken in 24 hours.

SIDE EFFECTS

When taken as directed, unwanted effects from Uphamol 650 are rare and usually mild. Skin rashes and other hypersensitivity reactions have been reported occasionally.

Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome/ Toxic Epidermal Necrolysis have been reported.

CONTRAINDICATION

Uphamol 650 should not be taken if you are allergic to acetaminophen.

WARNING/PRECAUTION

Please consult your doctor or pharmacist if pain persists.

Allergy alert: Paracetamol may cause severe skin reactions.

Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition.

If these reactions occur, stop use and seek medical assistance right away.

This preparation contains PARACETAMOL.
Do not take any other PARACETAMOL - containing medicines at the same time.

INTERACTIONS WITH OTHER MEDICAMENTS

Drugs which induce the specific microsomal isoenzyme responsible from the metabolic activation of paracetamol might be expected to increase susceptibility to its toxic effects on the liver humans as in the case in animals.

The absorption of paracetamol may accelerated by drugs such as metoclopramide. Excretion may be affected and plasma concentrations altered when given probenecid. Colestyramine reduces the absorption of paracetamol if given within 1 hour of paracetamol. Drug interaction also reported with warfarin, rifampicin, chloramphenicol, antiepileptic (carbamazepine, phenorbital, phenytoin, or primidone), antiviral (interferon alfa and zidovudine) and probenecid.

PREGNANCY AND LACTATION

Paracetamol crosses the placenta but the drug has been used widely as an analgesic in pregnancy and no adverse fetal affects has been recorded.