HALEON

Panadol C

Paracetamol 500mg • Pseudoephedrine Hydrochloride 30mg

Product Description
A White, capsule-shaped tablet with flat edges with face marked 'Panadol C' consist of paracetamol 500mg and pseudoephedrine hydrochloride 30mg.

Pharmacodynamic
Paracetamol
Paracetamol is an analgesic and antipyretic. Its mechanism of action is believed to include inhibition of prostaglandin synthesis, primarily within the central nervous system.
The lack of peripheral prostaglandin inhibition confers important pharmacological properties such as the maintenance of the protective prostaglandins within the gastrointestinal tract. Paracetamol is, therefore, particularly suitable for patients with a history of disease or on concomitant medication, where peripheral prostaglandin inhibition would be undesirable (such as, for example, those with a history of gastrointestinal bleeding or the elderly).

Pseudoephedrine Hydrochloride
Pseudoephedrine is predominantly an indirect-acting sympathomimetic amine with less direct adrenergic effect than its epimer, ephedrine. S(+)-pseudoephedrine is the naturally-occurring isomer and the one used in clinical practice.
Pseudoephedrine 60 mg has been shown to be an effective nasal decongestant, as measured by nasal airflow, in patients with the common cold and rhinitis and following histamine challenge in normal subjects and in patients with allergic rhinitis.
At therapeutic doses, pseudoephedrine has no clinically significant effect on blood pressure in normal subjects and patients with mild to moderate hypertension.

Pharmacokinetics
Paracetamol
Paracetamol is rapidly and almost completely absorbed from the gastrointestinal tract.
Binding to the plasma proteins is minimal at therapeutic concentrations.
Paracetamol is metabolised in the liver and excreted in the urine mainly as glucuronide and sulphate conjugates.
Less than 5% is excreted as unmodified paracetamol.

Pseudoephedrine Hydrochloride
Pseudoephedrine is rapidly and completely absorbed from the gastrointestinal tract after oral administration, with no presystemic metabolism.
Peak plasma levels are achieved after 1-2 hours. The plasma half-life varies from 4.3-7.0 hours in adults. No protein binding data are available.
There is little metabolism of pseudoephedrine in man with approximately 90% being excreted in the urine unchanged.
Approximately 1% is eliminated by hepatic metabolism, by N-demethylation to norpseudoephedrine
As a weak base, the extent of renal excretion is dependent on urinary pH. At low urinary pH, tubular reabsorption is minimal and urine flow rate will not influence clearance of the drug. At high pH (>7.0), pseudoephedrine is extensively reabsorbed in the renal tubule and renal clearance will depend on urine flow rate.
Renal impairment will increase plasma levels of pseudoephedrine.

What is this for?
For the temporary relief of sinus congestion and pain associated with rhinitis or the common cold.

How to use?
For oral administration.
Do not take more frequently than every four hours.
Do not exceed stated dose
Do not use with other decongestant or paracetamol-containing products.
Consult a doctor if symptoms persist for more than 7 days.
Not suitable for children under 12 years of age.
Tablets should be taken with water.
Adults (including the elderly) and children 12 years and over: One to two tablets every four to six hours, to be taken orally. Do not use more than 8 tablets in 24 hours.

Contraindications
Do not use if:
- You are allergic to paracetamol, pseudoephedrine or any of the other ingredients
- You have high blood pressure or heart disease
- You are taking other medicines for the relief of colds and flu, congestion or blocked nose, stimulant drugs called amphetamines (sometimes used to treat attention deficit disorders), or appetite suppressants
- You are taking or have taken in the past two weeks, drugs called monoamine oxidase inhibitors (MAOI's)
- You have kidney problems, unless your doctor tells you to

Warnings and Precautions
Check with your doctor before use if:
- You have liver problems
- You have an overactive thyroid gland, prostrate gland enlargement, an irregular heartbeat, glaucoma (excessive pressure inside your eyes), diabetes or phaeochromocytoma
- You are taking medicines to control your blood pressure e.g. medicines called alpha or beta blockers.
- If you are pregnant or breastfeeding
- You are taking warfarin or similar medicines used to thin the blood

Stop using this product and seek immediate medical attention if you experience sudden severe headache.
Users should be aware that this product can cause dizziness as side effect which could affect your ability to drive or operate machinery.
Please see your doctor if your symptoms do not improve. Keep out of sight and reach of children.

NPA018577

Interactions
The anticoagulant effect of warfarin and other coumarins may be enhanced by prolonged regular daily use of paracetamol-containing products with increased risk of bleeding; occasional doses have no significant effect.
Concomitant administration of pseudoephedrine hydrochloride-containing products and MAOIs (or within two weeks of stopping MAOI) may lead to hypertensive crisis.
Concomitant use of this medication with sympathomimetic agents (such as decongestant, appetite suppressant and amphetamine-like psychostimulants) which interfere with the catabolism of sympathomimetic amines may occasionally cause a rise in blood pressure.
Pseudoephedrine-containing products may antagonize the effect of certain classes of antihypertensive (e.g. beta blockers, methyl-dopa, reserpine, debrisoquine, guanethidine).

Pregnancy and Lactation
Pregnancy
This product should not be used in pregnancy without medical advice.
Safe use of pseudoephedrine in pregnancy has not been established despite widespread use over many years. Caution should therefore be exercised by balancing the potential benefit of treatment to the mother against any possible hazards to the developing foetus.
Lactation
This product should not be used whilst breastfeeding without medical advice.
Pseudoephedrine is excreted in breast milk in small amounts but the effect of this on breastfed infants is unknown.

Side Effect/Adverse Reactions
When using this product you may experience:
Nervousness, difficulty sleeping, dizziness, dry mouth, nausea and vomiting may occasionally occur. Hallucinations have been reported rarely.
Stop using this product and consult your doctor immediately if:
- You have difficulty passing water. This is more likely to occur if you have an enlarged prostrate gland.
- You experience allergic reactions such as skin rash or itching, sometimes with breathing problems or swelling of the lips, tongue, throat or face.
- You experience a skin rash or peeling, or mouth ulcers.
- You have previously experienced breathing problems with aspirin or non-steroidal anti-inflammatories, and experience a similar reaction with this product.
- You experience unexplained bruising or bleeding.

Overdose and Treatment
If you take more of the medicine than you should:
Seek medical advice immediately because of the risk of liver failure.

Paracetamol
Sign and symptoms
Paracetamol overdose, If you take more may cause the risk of liver failure.
Treatment
Immediate medical management is required in the event of overdose, even if symptoms of overdose are not present.
Administration of N-acetylcysteine or methionine may be required.

Pseudoephedrine Hydrochloride
Sign and symptoms
Pseudoephedrine overdose may result in symptoms due to central nervous system and cardiovascular stimulation e.g. excitement, restlessness, hallucinations, hypertension and arrhythmias. In severe cases, psychosis, convulsion, coma and hypertensive crisis may occur. Serum potassium levels may be low due to extracellular to intracellular shifts in potassium.
Treatment
Treatment should consist of standard supportive measures. Beta blockers should reverse the cardiovascular complication and the hypokalemia.

Pharmaceutical Information

Pack size
Blister of 10's
Box of 20 blisters of 10's.

How to store?
Store below 30°C.

Expiry
Do not use beyond the expiry date.

Manufactured by: Sterling Drug (Malaya) Sdn. Bhd. (4578-P)
Lot 89, Jalan Enggang, Ampang/Hulu Kelang Industrial Estate, 68000 Ampang, Selangor, Malaysia.
For: Haleon Malaysia Sdn. Bhd. [Registration No: 195901000115 (3467-X)]
Lot 89, Jalan Enggang, Ampang/Hulu Kelang Industrial Estate, 68000 Ampang, Selangor, Malaysia.
DATE OF REVISION: November 2025

mystory.my@haleon.com 1800 88 3225

Trade Marks are owned by or licensed to the Haleon group of companies.
©2025 Haleon group of companies or its licensor.

NPA018577