

LINCOMYCIN

 Injection "Yung Shin"

Lincomycin Injection contains Lincomycin hydrochloride which is the monohydrated salt of Lincomycin, a substance produced by the growth of a member of the *Lincolnensis* group of *Streptomyces lincolnensis*.

Ingredient: Each ml contains:

Lincomycin hydrochloride equivalent to Lincomycin 300mg (Potency)

Actions:

1. Lincomycin is effective against most of the common gram-positive pathogens. Depending on the sensitivity of the organism and concentration of the antibiotic, it may be either bactericidal or bacteriostatic.
2. Cross resistance has not been demonstrated with Penicillin, Chloramphenicol, Ampicillin, Cephalosporins or the Tetracyclines. Despite chemical differences, Lincomycin exhibits antibacterial activity similar but not identical to the Macrolide antibiotics (e.g. Erythromycin). Some cross resistance (with Erythromycin) including a phenomenon known as dissociated cross resistance or macrolide effect has been reported.
3. Microorganisms have not developed resistance to Lincomycin rapidly when tested by in vitro or in vivo methods.
4. In vitro studies indicate that the spectrum of activity includes *Staphylococcus aureus*, *Staphylococcus albus*, β -hemolytic *Streptococcus*, *Streptococcus viridans*, *Diplococcus pneumoniae*, *Clostridium tetani*, *Clostridium perfringens*, *Corynebacterium diphtheriae* and *Corynebacterium acnes*.
5. Intramuscular administration of a single dose of 600mg produces a peak serum level at 30 minutes with detectable levels persisting for 24 hours. Urinary excretion after this dose ranges from 1.8 to 24.8 percent (mean: 17.3%).
6. The intravenous infusion over a 2-hour interval of 600mg of Lincomycin potency in 500ml of 5% Glucose in distilled water yields therapeutic levels for 14 hours. Urinary excretion ranges from 4.9 to 30.3 percent. (mean: 13.8%).
7. The biological half-life, after intravenous or intramuscular administration is 5.4 ± 1.0 hours. Hemodialysis and peritoneal dialysis do not effectively remove Lincomycin from the blood.

Indications:

For the treatment of serious infections due to *Streptococci*, *Pneumococci*, *Staphylococci*, *Meningococci* and other susceptible organisms. It may be administered concomitantly with other antimicrobial agents when indicated.

Dosage and Administration:**Intramuscular:**

- Adults: Serious infections—600mg (2ml) intramuscularly every 24 hours.
More severe infections—600mg (2ml) intramuscularly every 12 hours or more often.
- Children over 1 month of age:
Serious infection— one intramuscular injection of 10mg/kg (5mg/lb) every 24 hours.
More severe infections— one intramuscular injection of 10mg/kg (5mg/lb) every 12 hours or more often.

Intravenous:

Adults: For serious infections doses of 600mg (2ml) to 1gm (3.3ml) are given every 8-12 hours. For more severe infections these doses may have to be increased. In life-threatening situations daily intravenous doses of as 8gm (26.7ml) have been given. Intravenous doses are given on the basis of 1gm of Lincomycin diluted in not less than 100ml of appropriate solution and infused over a period of not less than 1 hour.

Children over 1 month of age: 10-20mg/kg/day (5-10mg/lb/day) depending on the severity of the infection may be infused in divided doses as described above for adults. To be used only by a physician.

Warnings and Precautions:

1. Lincomycin therapy has been associated with severe colitis which may end fatally.
2. It should be reserved for serious infections where less toxic antimicrobial agents are inappropriate.
3. It should not be used in patients with nonbacterial infections, such as most upper respiratory tract infections.
4. Its use in newborns is contraindicated.
5. As this preparation contains benzyl alcohol, its use should be avoided in children under two years of age. Not to be used in neonates.

Packagings:

1mlx1vial, 1mlx10vial, 2mlx1vial, 2mlx10vial, 10mlx1vial, 10mlx10vial

Shelf-life:

3 years from the date of manufacture.

Storage Conditions:

Preserve in multiple-dose container, preferably of Type I glass.

Manufactured by:
YUNG SHIN PHARMACEUTICAL IND. CO., LTD.
NO. 1191, SEC. 1 CHUNG SHAN RD., TACHIA, TAICHUNG,
TAIWAN, R.O.C.

Imported & Distributed by:
Y.S.P. INDUSTRIES (M) SDN. BHD.
NO. 18, JALAN WAN KADIR, TAMAN TUN DR. ISMAIL,
60000, KUALA LUMPUR, MALAYSIA



2145642

