

PORO[®] TABLET

Ingredient(s):

Each tablet contains:

Paracetamol (Acetaminophen) 500mg

Action(s):

1. Paracetamol (Acetaminophen) may act by inhibiting prostaglandin synthesis in the central nervous system and through a peripheral action by blocking pain impulse generation. The peripheral action may also be due to inhibition of the synthesis of prostaglandins which sensitized pain receptors to mechanical or chemical stimulation.
2. Paracetamol produces antipyresis by acting centrally on the hypothalamic heat-regulation center to produce peripheral vasodilation resulting in increased blood flow through the skin, sweating and heat loss. The central action probably involves inhibition of prostaglandin synthesis in the hypothalamus.
3. Paracetamol is readily and completely absorbed from the gastrointestinal tract with peak plasma concentration occurring between 15 minutes to 2 hours after ingestion. Paracetamol is uniformly distributed throughout most body fluids; the apparent volume of distribution is 1 to 1.2 L/kg. Paracetamol may cross placenta and excreted in milk. At therapeutic plasma levels, no protein binding occurs.
Paracetamol is metabolized by the hepatic microsomal enzyme system. In adults at therapeutic doses, paracetamol is mainly conjugated with glucuronide (45-55%) or sulphate (20-30%). A minor proportion (less than 20%) is metabolized to catechol derivatives, and mercapturic acid compounds via oxidation. Paracetamol is metabolised differently by infants and children compared to adults, the sulphate conjugate being predominant.
Paracetamol is excreted by active tubular secretion: 85-90% of the administered dose is eliminated in the urine within 24 hours of administration. Paracetamol has a half-life of 1 to 4 hours. Less than 5 % is excreted as unchanged paracetamol.

Indication(s):

For the symptomatic relief of mild to moderate pain and to reduce fever.

Dosage and Administration:

The usual adult dose is 1 - 2 tablets 3 - 4 times daily. (Not more than 8 tablets / day unless prescribed by a physician). Do not exceed the prescribed dose. If symptoms persist, consult your doctor.

Route of Administration:

To be taken orally

Contraindication(s):

Contraindicated in patients who have shown hypersensitivity to Paracetamol.

Precaution(s) / Warning(s):

Paracetamol should be used with caution to patients with hepatic or renal dysfunction.

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash.

These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Side Effect(s) / Adverse Reaction(s):

Reports of adverse reactions are rare. Although dyspepsia, nausea, allergic and hematological reactions have been reported, a causal relationship to the administration of Paracetamol has neither been confirmed nor refuted.

Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome/Toxic Epidermal Necrolysis have been reported.

Symptoms and Treatment for Overdosage, and Antidote(s):

Toxic symptoms include vomiting, abdominal pain, hypotension and sweating. The most serious adverse effect of acute overdosage of Paracetamol is dose dependent, potentially fatal hepatic necrosis. In adults, hepatotoxicity may occur after ingestion of a single dose of Paracetamol 10-15g; a dose of 25g or more is potentially fatal.

In case of overdosage, methods of reducing the absorption of ingested drug are important. Gastric lavage is important even if several hours have elapsed. Prompt administration of activated charcoal 50g and 500ml of iced Mannitol 20% by mouth may reduce absorption.

If the history suggests that 15g or more has been ingested, administer one of the antidotes as follows: intravenous Acetylcystein 20%, oral Methionine and oral Acetylcysteine 5%.

Storage Condition(s):

Keep in a tight container. Store at temperature below 30°C. Protect from light and moisture.

Shelf-Life:

3 years from the date of manufacture.

Product Description and Packing(s):

Yellow color tablet, one side impressed with a score “” and the other side impressed with a mark “”.

Plastic bottle of 500's, 1000's, 1500's and 3000's (for export only)

Blister packing of 10's x 2, 10's x 3, 10's x 10, 10's x 100 and 10's x 300.

This preparation contains PARACETAMOL. Do not take any other paracetamol-containing medicines at the same time.

PORO[®] TABLET

Bahan aktif:

Setiap tablet mengandungi:

Paracetamol (Acetaminophen) 500mg

Indikasi:

Untuk melegakan kesakitan yang ringan dan sederhana serta mengurangkan demam.

Dos:

Dewasa: 1 – 2 tablet, 3 – 4 kali sehari. (Jangan mengambil melebihi 8 tablet / hari melainkan dipreskrib oleh doktor).

藥效確實、迅速、安全的鎮痛/解熱劑

伯樂錠

成份:

每錠含:

Paracetamol (Acetaminophen) 500mg

適應症:

- 伯樂錠是一種溫和的止痛劑，能夠提昇疼痛的閾值。本劑適用於緩解下列症狀所引起的疼痛：
 - 頭痛
 - 神經痛
 - 肌肉疼痛
 - 關節痛
 - 月經痛
- 本劑亦作用在下視丘之體溫控制中樞，促進皮膚層的血流量、造成出汗及解熱。故伯樂錠也適用於感冒、發燒的病症。
- 本劑口服後可迅速地由胃腸道吸收，於 15 分鐘到 2 個小時內發生作用，其藥效可維持 1 ~ 4 小時之久。

用法.用量:

成人: 每次1 ~ 2 粒, 每天3 ~ 4 次, 最高劑量為每天 8 粒, 或依醫師指示服用。



Y. S. P.

Manufacturer and Product Registration Holder:

Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)

Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.

Ordering Line: 1 800 88 3027

Product Info: 1 800 88 3679