

200mm

Front

148.5mm

pharmaniaga®

Isoniazid

Tablet 400mg

DESCRIPTION
Yellow, round, flat bevelled edge tablet with KK marking on one side and quadrisection on the other.

COMPOSITION
Each tablet contains 400 mg of isoniazid.

PHARMACODYNAMIC
Isoniazid has no significant antibacterial action against any micro-organisms except the mycobacteria; against mycobacterium tuberculosis it is bacteriostatic in extremely low concentrations.

Isoniazid is used mainly in the treatment of pulmonary tuberculosis but it appears to be effective also in the treatment of extrapulmonary lesions, including meningitis and genito-urinary disease.

PHARMACOKINETICS
Absorption: Readily and completely absorbed after oral administration.

Distribution: Readily diffuses into all tissues and fluids including the cerebrospinal fluid. Isoniazid is retained in the skin and in infected tissue; it crosses the placenta and is secreted in the milk of lactating mothers.

Protein binding: Isoniazid does not appear to be bound in the blood.

Half-life: Plasma elimination half-life, in rapid acetylators about 1.2 hours and in slow acetylators about 3.5 hours.

Metabolic reactions: Acetylation, hydrolysis and glycine conjugation, hydrazone formation, and n-methylation; acetylation is polymorphic and two groups of acetylators have been identified, rapid and slow acetylators. The rate of hydrolysis is more rapid in the rapid acetylators than in the slow ones. The metabolites formed include acetyl isoniazid, isonicotinic acid, isonicotinuric acid, isonicotinoyl-hydrazones of pyruvic and glutaric acids, and n-methylisoniazid.

Excretion: Over 90% of a dose is excreted in the urine in 24 hours, most being excreted in the first 12 hours, 4-32% is unchanged, but no more than 10% of a dose is excreted in the faeces.

INDICATIONS
Isoniazid is used primarily in conjunction with other anti-tuberculosis drugs in the treatment of all forms of tuberculosis. Isoniazid is also used in the treatment of tuberculosis meningitis and non-tuberculosis mycobacterium infections.

CONTRAINDICATIONS
Patients who are known to be hypersensitive to isoniazid or drug-induced liver disease.

ADVERSE REACTIONS
Blood and lymphatic system disorders
Frequency not known: Agranulocytosis, Aplastic anaemia, Haemolytic anaemia

Ear and labyrinth disorders
Frequency not known: Deafness, Tinnitus, Vertigo. These have been reported in patients with end stage renal impairment. Vertigo may be troublesome with doses of 10mg per kg body weight

Gastrointestinal disorders
Frequency not known: Constipation, Dry mouth Nausea, Pancreatitis acute, Vomiting and other gastrointestinal effects

General disorders and administration site conditions
Frequency not known: Pyrexia

Hepatobiliary disorders
Frequency uncommon: Hepatitis
Frequency not known: Acute hepatic failure, Liver injury, Jaundice

The risk of these undesirable effects increases with age, especially over the age of 35; it may be serious and sometimes fatal with the development of necrosis.

Investigations
Frequency not known: Hepatic enzyme increased

Metabolism and nutrition disorders
Frequency not known: Acidosis, Hypoglycaemia, Nicotinic acid deficiency. Nicotinic acid deficiency may be related to an isoniazid-induced pyridoxine deficiency which affects the conversion of tryptophan to nicotinic acid.

Musculoskeletal and connective tissue disorders
Frequency not known: Systemic lupus erythematosus, lupus-like syndrome

Nervous system disorders
Frequency not known: Neuropathy peripheral, Optic neuritis, Seizure. Hyperreflexia may be troublesome with doses of 10mg per kg body weight

Psychiatric disorders
Frequency not known: Elevated mood, Psychotic disorder.

Although isoniazid usually has a mood elevating effect, mental disturbances, ranging from minor personality changes to major mental derangement have been reported; these are usually reversed on withdrawal of the drug

Renal and urinary disorders
Frequency not known: Dysuria

Reproductive system and breast disorders
Frequency not known: Gynaecomastia

Respiratory, thoracic and mediastinal disorders
Frequency not known: Interstitial lung disease

Skin and subcutaneous tissue disorders
Frequency rare: Toxic epidermal necrolysis, eosinophilia systemic symptoms, Frequency not known: Erythema multiforme, Stevens-Johnson syndrome,

Vascular disorders
Frequency not known: Vasculitis

Miscellaneous
Withdrawal symptoms, which may occur on the cessation of the treatment, include headache, insomnia, excessive dreaming, irritability and nervousness

WARNINGS AND PRECAUTIONS
All patients should have baseline liver function tests performed and repeated at regular intervals during treatment. If serum AST rises to more than three times normal, or there is any increase in bilirubin, treatment should be withdrawn. Special precautions are required in patients with impaired liver function. Any deterioration in liver function in these patients is an indication for stopping treatment.


Isoniazid should not be given to patients who have experience severe adverse reactions including drug-induced liver disease. Care should be taken in giving isoniazid to patients suffering from convulsive disorders, diabetes mellitus, chronic alcoholism, or impaired liver or kidney function or to patients taking other potentially hepatotoxic agents. If symptoms of hepatitis such as malaise, fatigue, anorexia, and nausea develop isoniazid should be discontinued immediately.

Isoniazid should be used with caution in patients with a history of psychosis.

attn	customer Pharmaniaga Manufacturing Berhad	date 04.02.2021
1 colours	process colour: Black	Please Chop & Sign For Approval
size 200 x 148.5mm description Isoniazid Tablet 400mg - Front		date:
FocusPrint SDN BHD		
t: 603-8766 6030 f: 603-8766 6033 (Factory) website: www.focusprint.com.my		
artwork prepared by: cynthia yap		email: graphic@focusprint.info (graphic Dept)

IMPORTANT ! Please note that this colour print is for visual purpose only. Output colour might differ in actual production.

ARTWORK LOG		
Version no.	Date	Reason for Change
01	03.04.19	- Amend wording.
02	03.05.19	- Amend wording and date of revision.
03	13.05.19	- Amend date of revision to May.
04	13.05.19	- Amend PRP no.
05	24.05.19	- Amend TM to ® and date of revision.
06	07.10.2020	- Amend wording and PRP no.
07	26.11.2020	- Amend wording.
08	04.02.2021	- Amend wording.



attn		customer Pharmaniaga Manufacturing Berhad		date 04.02.2021
1 colours	process colour:	Black	Please Chop & Sign For Approval	
	size 200 x 148.5mm			
	description Isoniazid Tablet 400mg - Back			
 FocusPrint SDN BHD		t: 603-8766 6030 f: 603-8766 6033 (Factory) website: www.focusprint.com.my		date:
artwork prepared by: cynthia yap		email: graphic@focusprint.info (graphic Dept)		

IMPORTANT ! Please note that this colour print is for visual purpose only. Output colour might differ in actual production.

ARTWORK LOG		
Version no.	Date	Reason for Change
01	03.04.19	- Amend wording.
02	03.05.19	- Amend wording and date of revision.
03	13.05.19	- Amend date of revision to May.
04	13.05.19	- Amend PRP no.
05	24.05.19	- Amend TM to ® and date of revision.
06	07.10.2020	- Amend wording and PRP no.
07	26.11.2020	- Amend wording.
08	04.02.2021	- Amend wording.