

Important Information. Please read carefully.

AVORAX[®]

Cream



Composition

Contains Acyclovir 5% w/w in a water miscible cream.
Chlorocresol 0.10% w/w as preservative.

Pharmacodynamics

Acyclovir is an antiviral agent which is highly active in vitro against herpes simplex virus (HSV) types I and II and varicella zoster virus. Toxicity to mammalian host cells is low. Acyclovir is phosphorylated after entry into herpes infected cells to the active compound acyclovir triphosphate. The first step in this process is dependent on the presence of the HSV-coded thymidine kinase. Acyclovir triphosphate acts as an inhibitor of, and substrate for, the herpes-specified DNA polymerase, preventing further viral DNA synthesis without affecting normal cellular processes.

Pharmacokinetics

Pharmacology studies have shown only minimal systemic absorption of acyclovir following repeated topical administration of Avorax Cream.

Indications

Avorax Cream is indicated in the treatment of herpes simplex infections of the skin, including initial and recurrent labial and genital herpes simplex infections.

Dosage and Administration

For topical use only. Avorax Cream should be applied every 3 - 4 hours or 5 times daily for 5 - 10 days. Avorax should be applied as soon as infection is detected and treatment must proceed for another 5 days after healing.

Contraindications

It is contraindicated in patients with known hypersensitivity to Acyclovir.

Precautions

Acyclovir Cream is not recommended for use on mucous membranes such as in the mouth, eye or vagina. Although no information is yet available on the adverse effects of use during pregnancy and nursing mothers, care should be taken during usage in such conditions.

Drug Interactions

No clinically significant interactions have been identified.

Incompatibilities

None known.

Usage during Pregnancy

The use of Avorax Cream should be considered only when the potential benefits outweigh the possibility of unknown risks however the systemic exposure to acyclovir from topical application of Avorax Cream is very low.

Overdosage and Treatment

No untoward effects would be expected if the acyclovir cream were ingested orally. However the accidental, repeated overdose of oral acyclovir, over several days has resulted in gastrointestinal effects (nausea and vomiting) and neurological effects (headache and confusion). Acyclovir is dialysable by haemodialysis.

Side Effects

Side effects includes transient stinging or burning sensation on application. Few patients have also reported erythema or drying of the skin.

Presentation

In collapsible aluminium tubes 5gm.

Shelf-Life

The expiry date is indicated on the packaging.

Storage

Protect from light. Store in a cool, dry place below 30°C. Keep tube well closed.

**KEEP OUT REACH OF CHILDREN
JAUHI DARI KANAK-KANAK**

For further information, please consult your pharmacist or physician.

Revision Date : 26-Oct-2020

Manufacturer and Product Registration Holder
Xepa-Soul Pattinson (Malaysia) Sdn Bhd
1-5 Cheng Industrial Estate, 75250 Melaka, Malaysia.