
ANALPAN

DESCRIPTION:

ANALPAN INJECTION 75mg / 3ml: A clear, colourless to almost colourless liquid in ampoule for IV infusion or I/M injection.
ANALPAN INJECTION 25mg / ml: A clear, colourless to almost colourless liquid in vial for IV infusion or I/M injection.

COMPOSITION:

Each ml contains 25mg/ml Diclofenac Sodium.

PHARMACODYNAMICS:

Analpan Injection contains diclofenac sodium, a non-steroidal compound with pronounced antirheumatic, anti-inflammatory, analgesic, and antipyretic properties. Inhibition of prostaglandin biosynthesis, which has been demonstrated in experiments, is considered to be fundamental to its mechanism of action. Prostaglandins play an important role in causing inflammation, pain and fever. Diclofenac sodium in vitro does not suppress proteoglycan biosynthesis in cartilage at concentrations equivalent to the concentrations reached in humans.

In rheumatic diseases, the anti-inflammatory and analgesic properties of diclofenac elicit a clinical response characterized by marked relief from signs and symptoms such as pain at rest, pain on movement, morning stiffness, and swelling of the joints, as well as by an improvement in function.

In post-traumatic and post-operative inflammatory conditions, diclofenac rapidly relieves both spontaneous pain and pain on movement and reduces inflammatory swelling and wound edema.

Analpan Injection has also been found to exert a pronounced analgesic effect in moderate and severe pain of non-rheumatic origin, an effect which sets in within 15 to 30 minutes.

When used concomitantly with opioids for the management of post-operative pain, Analpan Injection significantly reduces the need for opioids. Analpan Injection are particularly suitable for initial treatment of inflammatory and degenerative rheumatic diseases, and of painful conditions due to inflammation of nonrheumatic origin.

PHARMACOKINETICS:

Absorption:

After administration of 75 mg diclofenac by intramuscular injection, absorption sets in immediately, and mean peak plasma concentrations of about 2.5 micrograms/mL (8micromol/L) are reached after about 20 minutes.

When 75 mg diclofenac is administered as an intravenous infusion over 2 hours, mean peak plasma concentrations are about 1.9 micrograms/mL (5.9 micromol/L). Shorter infusions result in higher peak plasma concentrations, while longer infusions give plateau concentrations proportional to the infusion rate after 3 to 4 hours. In contrast, plasma concentrations decline rapidly once peak levels have been reached following intramuscular injection or administration of gastro-resistant tablets or suppositories.

The area under the concentration curve (AUC) after intramuscular or intravenous administration is about twice as large as it is following oral or rectal administration, because about half the active substance is metabolized during its first passage through the liver ("first pass" effect) when administered via the oral or rectal routes.

Pharmacokinetic behaviour does not change after repeated administration. No accumulation occurs provided the recommended dosage intervals are observed.

Distribution:

99.7% of diclofenac binds to serum proteins, mainly to albumin (99.4%). The apparent volume of distribution calculated is 0.12 to 0.17 L/kg.

Diclofenac enters the synovial fluid, where maximum concentrations are measured 2 to 4 hours after peak plasma values have been reached. The apparent half-life for elimination from the synovial fluid is 3 to 6 hours. Two hours after reaching peak plasma levels, concentrations of the active substance are already higher in the synovial fluid than in the plasma, and they remain higher for up to 12 hours.

Biotransformation/Metabolism: Biotransformation of diclofenac takes place partly by glucuronidation of the intact molecule, but mainly by single and multiple hydroxylation and methoxylation, resulting in several phenolic metabolites (3'-hydroxy-, 4'-hydroxy-, 5-hydroxy-, 4',5-dihydroxy-, and 3'-hydroxy-4'-methoxy-diclofenac), most of which are converted to glucuronide conjugates. Two of these phenolic metabolites are biologically active, but to a much lesser extent than diclofenac.

Elimination:

Total systemic clearance of diclofenac from plasma is 263 ± 56 mL/min (mean value $\pm$ SD). The terminal half-life in plasma is 1 to 2 hours. Four of the metabolites, including the two active ones, also have short plasma half-lives of 1 to 3 hours. One metabolite, 3'-hydroxy-4' methoxydiclofenac, has a much longer plasma half-life. However, this metabolite is virtually inactive.

About 60% of the administered dose is excreted in the urine as the glucuronide conjugate of the intact molecule and as metabolites, most of which are also converted to glucuronide conjugates. Less than 1% is excreted as unchanged substance. The rest of the dose is eliminated as metabolites through the bile in the faeces.

INDICATIONS:

Intramuscular injection

Treatment of:

- Exacerbations of inflammatory and degenerative forms of rheumatism: rheumatoid arthritis, ankylosing spondylitis, osteoarthritis, spondylarthritis, painful syndromes of the vertebral column, non-articular rheumatism.
- Acute attacks of gout.
- Renal colic and biliary colic.
- Post-traumatic and post-operative pain, inflammation and swelling.

Intravenous infusion

Treatment or prevention of post-operative pain in a hospital setting

RECOMMENDED DOSAGE:

Dosage regimen

As a general recommendation, the dose should be individually adjusted. Adverse effects may be minimized by using the lowest effective dose for the shortest duration necessary to control symptoms (see section WARNINGS AND PRECAUTIONS).

General target population

Adults

Analpan injection should not be given for more than 2 days (see Method of administration).

Special populations

Renal impairment

Analpan Injection is contraindicated in patients with renal failure (GFR <15 mL/min/1.73 m²) (see section CONTRAINDICATIONS).

No specific studies have been carried out in patients with renal impairment, therefore, no specific dose adjustment recommendations can be made. Caution is advised when administering Analpan to patients with renal impairment (see section WARNINGS AND PRECAUTIONS).

Hepatic impairment

Analpan is contraindicated in patients with hepatic failure (see section CONTRAINDICATIONS).

No specific studies have been carried out in patients with hepatic impairment, therefore, no specific dose adjustment recommendations can be made. Caution is advised when administering Analpan to patients with mild to moderate hepatic impairment (see section WARNINGS AND PRECAUTIONS).

Pediatric patients (below 18 years)

Because of their dosage strength, the Analpan injection are not suitable for children and adolescents.

Geriatric patients (65 years of age or above)

No adjustment of the starting dose is generally required for elderly patients. However, caution is indicated on basic medical grounds, especially for frail elderly patients or those with a low body weight (see section WARNINGS

AND PRECAUTIONS).

Established cardiovascular disease or significant cardiovascular risk factors

Treatment with Analpan is generally not recommended in patients with established cardiovascular disease (congestive heart failure, established ischemic heart disease, peripheral arterial disease) or uncontrolled hypertension. If needed, patients with established cardiovascular disease, uncontrolled hypertension, or significant risk factors for cardiovascular disease (e.g. hypertension, hyperlipidaemia, diabetes mellitus and smoking) should be treated with Analpan only after careful consideration and only at doses ≤100 mg daily if treated for more than 4 weeks (see section WARNINGS AND PRECAUTIONS).

Method of administration

Intramuscular injection

The following directions for intramuscular injection must be followed in order to avoid damage to a nerve or other tissue at the injection site (which may result in muscle weakness, muscle paralysis, hypoesthesia and Embolia cutis medicamentosa (Nicolau syndrome)).

The dose is generally one 75 mg daily, given by deep intragluteal injection into the upper outer quadrant using aseptic technique. In severe cases (e.g., colic), the daily dose can exceptionally be increased to two injections of 75 mg, separated by an interval of a few hours (one into each buttock).

Intravenous infusion

Analpan injection must not be given as an intravenous bolus injection.

Immediately before starting an intravenous infusion, Analpan injection must be diluted with saline 0.9% or glucose 5% infusion solution buffered with sodium bicarbonate according to the instructions given in section INSTRUCTIONS FOR USE AND HANDLING.

Two alternative dosage regimens of Analpan injection are recommended.

For the treatment of moderate to severe post-operative pain, 75 mg should be infused continuously over a period of 30 minutes to 2 hours. If necessary, treatment may be repeated after a few hours, but the dose should not exceed 150 mg within any period of 24 hours.

For the prevention of post-operative pain, a loading dose of 25 mg to 50 mg should be infused after surgery over 15 minutes to 1 hour, followed by a continuous infusion of about 5 mg per hour up to a maximum daily dose of 150 mg.

INSTRUCTIONS FOR USE AND HANDLING

The following directions for intramuscular injection must be followed in order to avoid damage to a nerve or other tissue at the injection site.

To be injected either intramuscularly by deep intragluteal injection into the upper outer quadrant using aseptic technique, or intravenously by slow infusion after dilution in accordance with the following instructions. Each ampoule is for single use only. The solution should be used immediately after opening. Any unused contents should be discarded.

Appropriate injection technique and length of the needle (considering the thickness of the patient's gluteal fat) should be used to avoid inadvertent subcutaneous administration of Analpan injection.

Depending on the intended duration of infusion (see section DOSAGE REGIMEN AND ADMINISTRATION), mix 100 to 500 mL of isotonic saline (sodium chloride 0.9% solution) or glucose 5% solution with the contents of one Analpan injection. Both solutions should be buffered with sodium bicarbonate injectable solution (0.5 mL of 8.4% or 1 mL of 4.2% or a corresponding volume of a different concentration) taken from a freshly opened container. Only clear solutions should be used. If crystals or precipitates are observed, the infusion solution should not be used.

Each Analpan Injection ampoule (75mg/3ml) is for single use only. The solution should be used immediately after opening. Any unused contents should be discarded. Analpan Injection 25mg/ml (10ml vial) remained stable for up to 28 days at 30oC/75% RH after opened and withdrawn several times within 28 days.

ROUTE OF ADMINISTRATION: For Intravenous Infusion or Intramuscular (I.M) Injection.

CONTRAINDICATIONS:

- Known hypersensitivity to the active substance, sodium metabisulphite or to any of the other excipients.
- Active gastric or intestinal ulcer, bleeding or perforation (see sections WARNINGS AND PRECAUTIONS and ADVERSE DRUG REACTIONS).
- Last trimester of pregnancy (see section PREGNANCY, LACTATION, FEMALES AND MALES OF REPRODUCTIVE POTENTIAL).
- Hepatic failure.
- Renal failure (GFR <15 mL/min/1.73 m²).
- Severe cardiac failure (see section WARNINGS AND PRECAUTIONS).
- Like other non-steroidal anti-inflammatory drugs (NSAIDs), Analpan is also contraindicated in patients in whom the use of acetylsalicylic acid or other NSAIDs can precipitate asthma, angioedema, urticaria, or acute rhinitis (i.e., NSAID-induced cross reactivity reactions) (see sections WARNINGS AND PRECAUTIONS and ADVERSE DRUG REACTIONS).

WARNINGS & PRECAUTIONS:

Gastrointestinal effects

Gastrointestinal bleeding ulceration or perforation, which can be fatal, have been reported with all NSAIDs, including diclofenac, and may occur at any time during treatment, with or without warning symptoms or a previous history of serious gastrointestinal events. They generally have more serious consequences in the elderly. If gastrointestinal bleeding or ulceration occurs in patients receiving Analpan, the treatment should be discontinued.

As with all NSAIDs, including diclofenac, close medical surveillance is imperative and particular caution should be exercised when prescribing Analpan in patients with symptoms indicative of gastrointestinal (GI) disorders or with a history suggestive of gastric or intestinal ulceration, bleeding or perforation, (see section ADVERSE DRUG REACTIONS). The risk of GI bleeding is higher with increasing NSAID doses and in patients with a history of ulcer, particularly if complicated with hemorrhage or perforation and in the elderly.

To reduce the risk of GI toxicity in patients with a history of ulcer, particularly if complicated with hemorrhage or perforation, and in the elderly, the treatment should be initiated and maintained at the lowest effective dose.

Combination therapy with protective agents (e.g., proton pump inhibitors or misoprostol) should be considered for these patients, and also for patients requiring concomitant use of low dose acetylsalicylic acid (ASA), or other drugs likely to increase gastrointestinal risk.

Patients with a history of GI toxicity, particularly the elderly, should report any unusual abdominal symptoms (especially GI bleeding). Caution is recommended in patients receiving concomitant medications which could increase the risk of ulceration or bleeding, such as systemic corticosteroids, anticoagulants, anti-platelet agents or selective serotonin-reuptake inhibitors (see section INTERACTIONS).

Close medical surveillance and caution should also be exercised in patients with ulcerative colitis or Crohn's disease, as their condition may be exacerbated (see section ADVERSE DRUG REACTIONS). NSAIDs, including diclofenac, may be associated with increased risk of gastro-intestinal anastomotic leak. Close medical surveillance and caution are recommended when using Analpan after gastro-intestinal surgery.

Cardiovascular effects

Treatment with NSAIDs including diclofenac, particularly at high dose and in long term, may be associated with an increased risk of serious cardiovascular thrombotic events (including myocardial infarction and stroke).

Treatment with Analpan is generally not recommended in patients with established cardiovascular disease (congestive heart failure, established ischemic heart disease, peripheral arterial disease) or uncontrolled hypertension. If needed, patients with established cardiovascular disease, uncontrolled hypertension, or significant risk factors for cardiovascular disease (e.g., hypertension, hyperlipidemia, diabetes mellitus and smoking) should be treated with Analpan only after careful consideration and only at doses ≤100 mg daily when treatment continues for more than 4 weeks.

As the cardiovascular risks of diclofenac may increase with dose and duration of exposure, the lowest effective daily dose should be used for the shortest duration possible. The patient's need for symptomatic relief and response to therapy should be re-evaluated periodically, especially when treatment continues for more than 4 weeks.

Patients should remain alert for the signs and symptoms of serious arteriothrombotic events (e.g., chest pain, shortness of breath, weakness, slurring of speech), which can occur without warnings. Patients should be instructed to see a physician immediately in case of such an event.

Hematologic effects

During prolonged treatment with Analpan, as with other NSAIDs, monitoring of the blood count is recommended. Like other NSAIDs, diclofenac may temporarily inhibit platelet aggregation. Patients with defects of hemostasis should be carefully monitored.

Respiratory effects (pre-existing asthma)

In patients with asthma, seasonal allergic rhinitis, swelling of the nasal mucosa (i.e., nasal polyps), chronic obstructive pulmonary diseases or chronic infections of the respiratory tract (especially if linked to allergic rhinitis-like symptoms), reactions on NSAIDs like asthma exacerbations (so-called intolerance to analgesics / analgesics-asthma), Quincke's edema or urticaria are more frequent than in other patients. Therefore, special caution is recommended in such patients (readiness for emergency). This is applicable as well for patients who are allergic to other substances, e.g., with skin reactions, pruritus or urticaria.

Special caution is recommended when Analpan is used parenterally in patients with bronchial asthma because symptoms may be exacerbated.

Hepatobiliary effects

Close medical surveillance is required when prescribing Analpan to patients with impaired hepatic function, as their condition may be exacerbated.

As with other NSAIDs, including diclofenac, values of one or more liver enzymes may increase. During prolonged treatment with Analpan (e.g., in the form of tablets or suppositories), regular monitoring of hepatic function is indicated as a precautionary measure. If abnormal liver function tests persist or worsen, if clinical signs or symptoms consistent with liver disease develop, or if other manifestations occur (e.g., eosinophilia, rash), Analpan should be discontinued. Hepatitis may occur with use of diclofenac without prodromal symptoms.

Caution is called for when using Analpan in patients with hepatic porphyria, since it may trigger an attack.

Skin reactions

Serious skin reactions, some of them fatal, including exfoliative dermatitis, Stevens-Johnson syndrome and toxic epidermal necrolysis, have been reported very rarely in association with the use of NSAIDs, including Analpan (see section ADVERSE DRUG REACTIONS). Patients appear to be at highest risk of these reactions early in the course of therapy, the onset of the reaction occurring in the majority of cases within the first month of treatment. Analpan should be discontinued at the first appearance of skin rash, mucosal lesions or any other sign of hypersensitivity.

As with other NSAIDs, allergic reactions, including anaphylactic/anaphylactoid reactions, can also occur in rare cases with diclofenac without earlier exposure to the drug.

Renal effects

As fluid retention and edema have been reported in association with NSAID therapy, including diclofenac, particular caution is called for in patients with impaired cardiac or renal function, history of hypertension, the elderly, patients receiving concomitant treatment with diuretics or medicinal products that can significantly impact renal function, and in those patients with substantial extracellular volume depletion of any cause, e.g., before or after major surgery (see section CONTRAINDICATIONS). Monitoring of renal function is recommended as a precautionary measure when using Analpan in such cases. Discontinuation of therapy is usually followed by recovery to the pre-treatment state.

Injection site reactions

Injection site reactions have been reported after the administration of Analpan intramuscularly, including injection site necrosis and embolia cutis medicamentosa, also known as Nicolau Syndrome (particularly after inadvertent subcutaneous administration). Appropriate needle selection and injection technique should be followed during i.m. administration of Analpan (see section INSTRUCTIONS FOR USE AND HANDLING).

Geriatric patients

Caution is indicated in the elderly on basic medical grounds, especially in frail elderly patients or those with a low body weight

Interactions with NSAIDs

The concomitant use of Analpan with systemic NSAIDs including cyclooxygenase-2 selective inhibitors, should be avoided due to undesirable effects (see section INTERACTIONS).

Special excipients

The sodium metabisulphite in the solution for injection can lead to isolated severe hypersensitivity reactions and bronchospasm.

Masking signs of infections

Like other NSAIDs, diclofenac may mask the signs and symptoms of infection due to its pharmacodynamic properties.

INTERACTION WITH OTHER MEDICATIONS:**Observed interactions to be considered**

CYP2C9 inhibitors: Caution is recommended when co-prescribing diclofenac with CYP2C9 inhibitors (such as voriconazole), which could result in a significant increase in peak plasma concentrations and exposure to diclofenac.

Lithium: If used concomitantly, diclofenac may raise plasma concentrations of lithium. Monitoring of the serum lithium level is recommended.

Digoxin: If used concomitantly, diclofenac may raise plasma concentrations of digoxin. Monitoring of the serum digoxin level is recommended.

Diuretics and antihypertensive agents: Like other NSAIDs, concomitant use of diclofenac with diuretics or antihypertensive agents (e.g., beta-blockers, angiotensin converting enzyme (ACE) inhibitors) may cause a decrease in their antihypertensive effect. Therefore, the combination should be administered with caution and patients, especially the elderly, should have their blood pressure periodically monitored. Patients should be adequately hydrated and consideration should be given to monitoring of renal function after initiation of concomitant therapy and periodically thereafter, particularly for diuretics and ACE inhibitors due to the increased risk of nephrotoxicity (see section WARNINGS AND PRECAUTIONS).

Ciclosporin and tacrolimus: Diclofenac, like other NSAIDs may increase the nephrotoxicity of ciclosporin and tacrolimus due to the effect on renal prostaglandins. Therefore, it should be given at doses lower than those that would be used in patients not receiving ciclosporin or tacrolimus.

Drugs known to cause hyperkalemia: Concomitant treatment with potassium-sparing diuretics, ciclosporin, tacrolimus or trimethoprim may be associated with increased serum potassium levels, which should therefore be monitored frequently (see section WARNINGS AND PRECAUTIONS).

Quinolone antibacterials: There have been isolated reports of convulsions which may have been due to concomitant use of quinolones and NSAIDs.

Anticipated interactions to be considered

Other NSAIDs and corticosteroids: Concomitant administration of diclofenac and other systemic NSAIDs or corticosteroids may increase the frequency of gastrointestinal undesirable effects (see section WARNINGS AND PRECAUTIONS).

Anticoagulants and anti-platelet agents: Caution is recommended since concomitant administration could increase the risk of bleeding (see section WARNINGS AND PRECAUTIONS). Although clinical investigations do not appear to indicate that diclofenac affects the action of anticoagulants, there are reports of an increased risk of hemorrhage in patients receiving diclofenac and anticoagulants concomitantly. Close monitoring of such patients is therefore recommended.

Selective serotonin reuptake inhibitors (SSRIs): Concomitant administration of systemic NSAIDs, including diclofenac, and SSRIs may increase the risk of gastrointestinal bleeding (see section WARNINGS AND PRECAUTIONS).

Antidiabetics: Clinical studies have shown that diclofenac can be given together with oral antidiabetic agents without influencing their clinical effect. However, there have been isolated reports of both hypoglycemic and hyperglycemic effects necessitating changes in the dosage of the antidiabetic agents during treatment with diclofenac. For this reason, monitoring of the blood glucose level is recommended as a precautionary measure during concomitant therapy. There have also been isolated reports of metabolic acidosis when diclofenac was coadministered with metformin, especially in patients with pre-existing renal impairment.

Phenytoin: When using phenytoin concomitantly with diclofenac, monitoring of phenytoin plasma concentrations is recommended due to an expected increase in exposure to phenytoin.

Methotrexate: Caution is recommended when NSAIDs, including diclofenac are administered less than 24 hours before or after treatment with methotrexate, since blood concentrations of methotrexate may rise and the toxicity of this substance be increased.

CYP2C9 inducers: Caution is recommended when co-prescribing diclofenac with CYP2C9 inducers (such as rifampicin), which could result in a significant decrease in plasma concentration and exposure to diclofenac.

USE IN PREGNANCY & LACTATION:**Pregnancy**

There are insufficient data on the use of diclofenac in pregnant women. Some epidemiological studies suggest an increased risk of miscarriage after use of a prostaglandin synthesis inhibitor (such as NSAIDs) in early pregnancy, however the overall data are inconclusive. Diclofenac has been shown to cross the placental barrier in humans. Use of NSAIDs, including diclofenac, can cause uterine inertia, premature closure of the fetal ductus arteriosus and fetal renal impairment leading to oligohydramnios. Because of these risks, Analpan should not be used during the first two trimesters of pregnancy unless the expected benefits to the mother outweigh the risks to the fetus. In addition, Analpan should not be used during the third trimester of pregnancy (see section CONTRAINDICATIONS).

In animal reproduction studies, no evidence of teratogenicity was observed in mice, rats, or rabbits given diclofenac daily during the period of organogenesis at doses up to approximately 0.41, 0.41, and 0.81 times, respectively, the maximum recommended human dose (MRHD) of Analpan, despite the presence of maternal and fetal toxicity.

Clinical considerations**Fetal Adverse Drug Reactions****Premature Closure of Fetal Ductus Arteriosus**

As with other NSAIDs, use of diclofenac during the third trimester of pregnancy is contraindicated owing to the possibility of premature closure of the fetal ductus arteriosus (see section CONTRAINDICATIONS).

Oligohydramnios/Fetal Renal Impairment

Risk of fetal renal impairment with subsequent oligohydramnios has been observed when NSAIDs (including diclofenac) were used from the 20th week of pregnancy onwards.

If an NSAID is necessary from the 20th week gestation to the end of the 2nd trimester, limit the use to the lowest effective dose and shortest duration possible (see section DOSAGE REGIMEN AND ADMINISTRATION).

If Analpan treatment extends beyond 48 hours, consider monitoring with ultrasound for oligohydramnios. If oligohydramnios occurs, discontinue Analpan and follow up according to clinical practice.

Labor or Delivery

There are no studies on the effects of Analpan during labor or delivery. As with other NSAIDs, use of diclofenac during the third trimester of pregnancy is contraindicated owing to the possibility of uterine inertia (see section CONTRAINDICATIONS). In animal studies, NSAIDs, including diclofenac, inhibit prostaglandin synthesis, cause delayed parturition, and increase the incidence of stillbirth.

Lactation

Like other NSAIDs, diclofenac passes into the breast milk in small amounts. Therefore, Analpan should not be administered during breast feeding in order to avoid undesirable effects in the infant.

Female fertility

As with other NSAIDs, the use of Analpan may impair female fertility and is not recommended in women attempting to conceive. In women who have difficulties conceiving or who are undergoing investigation of infertility, withdrawal of Analpan should be considered.

Male fertility

There is no human data on the effect of Analpan on male fertility.

Diclofenac administered to male and female rats at 4 mg/kg/day (approximately 0.16 times the MRHD based on BSA comparison) did not affect fertility.

SIDE EFFECTS/ ADVERSE EFFECTS:

Gastrointestinal Tract:

Common: Epigastric pain, other gastrointestinal disorders such as nausea, vomiting, diarrhoea, abdominal cramps, dyspepsia, flatulence, anorexia, decreased appetite.

Rare: Gastrointestinal bleeding, haematemesis, melaena, gastric or intestinal ulcer with or without bleeding or gastrointestinal stenosis, or perforation, which may lead to peritonitis, bloody diarrhoea. Gastritis

Very Rare: Lower gut disorders such as non-specific haemorrhagic colitis and exacerbation of ulcerative colitis or Crohn's proctocolitis; aphthous stomatitis, glossitis, oesophageal lesions, constipation, diaphragm-like intestinal strictures, pancreatitis.

Central (and peripheral) Nervous System:

Common: headache, dizziness, or vertigo.

Rare: Somnolence, tiredness.

Very Rare: disturbances of sensation, including paraesthesia, memory disturbance, disorientation, disturbance of vision (blurred vision, diplopia), impaired hearing; tinnitus, insomnia, irritability, convulsions, depression, anxiety, nightmares, tremor, psychotic reactions, taste alteration disorder, memory impairment, dysgeusia, cerebrovascular accident, myoclonic encephalopathy (described in two patients), aseptic meningitis.

Cardiac disorders:

Uncommon:* Myocardial infarction, cardiac failure, palpitations, chest pain. [*The frequency reflects data from long-term treatment with a high dose (150mg/day)]

Kounis syndrome: Frequency "not known"

Arteriothrombotic events:

Meta-analysis and pharmacoepidemiological data point towards an increased risk of arteriothrombotic events (for example myocardial infarction) associated with the use of diclofenac, particularly at a high dose (150mg daily) and during long-term treatment (see section WARNINGS AND PRECAUTIONS).

Dermatological:

Occasionally: rashes or skin eruptions, cases of hair loss, bullous eruptions, erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis (Lyell's syndrome), and photosensitivity reactions have been reported.

Rare: urticaria.

Very Rare: eczema, erythroderma (exfoliative dermatitis), purpura, including allergic purpura, Bullous dermatitis, erythema, erythema multiforme, pruritus.

Urogenital System: *Very Rare:* acute renal failure, urinary abnormalities such as haematuria, proteinuria, interstitial nephritis, nephrotic syndrome, papillary necrosis.

Liver:

Common: elevation of serum aminotransferase enzymes (ALT, AST)

Rare: hepatitis with or without jaundice, liver disorder.

Very Rare: fulminant hepatitis, hepatic necrosis, hepatic failure.

Blood: *Very Rare:* thrombocytopenia, leukopenia, anaemia (haemolytic anaemia, aplastic anaemia), agranulocytosis, positive Coombs' test.

Immune System Disorders:

Less than 1%: Hypersensitivity reactions such as bronchospasm, anaphylactic/anaphylactoid systemic reactions including hypotension. In isolated cases: Vasculitis, pneumonitis.

Very rare: Angioedema (including face edema).

Infections and infestations

Very rare: Injection site abscess.

General disorders and administration site conditions

Common: Injection site reaction, injection site pain, injection site induration.

Rare: Edema, injection site necrosis

Eye and Ear:

Common: Vertigo.

Very Rare: Visual disturbance, vision blurred, diplopia, Tinnitus, hearing impaired

Others: *In isolated cases:* impotence (association with diclofenac sodium intake is doubtful), palpitation, chest pain, hypertension, congestive heart failure.

SYMPTOMS AND TREATMENT OF OVERDOSE:

Symptoms

There is no typical clinical picture resulting from diclofenac overdosage. Overdosage can cause symptoms such as vomiting, gastrointestinal hemorrhage, diarrhea, dizziness, tinnitus or convulsions. In the event of significant poisoning, acute renal failure and liver damage are possible.

Therapeutic measures

Management of acute poisoning with NSAIDs, including diclofenac, essentially consists of supportive measures and symptomatic treatment. Supportive measures and symptomatic treatment should be given for complications such as hypotension, renal failure, convulsions, gastrointestinal disorder, and respiratory depression. Special measures such as forced diuresis, dialysis or hemoperfusion are probably of no help in eliminating NSAIDs, including diclofenac, due to the high protein binding and extensive metabolism.

INCOMPATIBILITIES:

As a rule, Analpan injection should not be mixed with other injection solutions. Infusion solutions of sodium chloride 0.9% or glucose 5% without sodium bicarbonate as an additive present a risk of supersaturation, possibly leading to formation of crystals or precipitates. Infusion solutions other than those recommended should not be used.

STORAGE CONDITIONS: Store below 30°C. Protect from light. Keep out of reach of children. Jauhkan daripada kanak-kanak

SHELF LIFE: Please refer to outer package.

PACK SIZE:

Ampoules : 3ml x 10 ampoules and 3ml x 100 ampoules.

Vials : 10ml x 25 vials

PRODUCT REGISTRATION HOLDER & MANUFACTURER:

DUOPHARMA (M) SDN BHD

Lot 2599 Jalan Seruling 59 Kawasan 3,

Taman Klang Jaya, 41200 Klang, Selangor, MALAYSIA