

Composition : Contains Ketoconazole 2%w/w in cream base.

Preservative : Methyl Hydroxybenzoate 0.2%w/w;

Propyl Hydroxybenzoate 0.1%w/w.

Properties : Ketoconazole is an imidazole antifungal agent that impairs the synthesis of ergosterol, which is a vital component of fungal cell membrane and therefore alters the permeability of the cell membrane of the following common dermatophytes and yeast. Dermatophytes : *Trichophyton rubrum*, *T. mentagrophytes*, *T. tonsurans*, *Microsporum canis*, *M. audouinii*, *M. gypseum* and *Epidermophyton floccosum*; Yeast: *Candida albicans*, *Malassezia ovale* (*Pityrosporum ovale*) and *C. tropicalis*; and the common organism responsible for tinea versicolor, *Malassezia furfur* (*Pityrosporum orbiculare*).

Indications : Ketoconazole cream is indicated for the topical treatment of tinea corporis (ringworm of the body), tinea cruris (jock itch), tinea pedis (athlete's foot), tinea (pityriasis) versicolor (white spots), cutaneous candidiasis and seborrheic dermatitis.

Dosage and Administration:

- ◆ For cutaneous candidiasis, ringworm of the body, jock itch, tinea manus, athlete's foot and tinea (pityriasis) versicolor (white spots):
- apply once daily to cover the affected and immediate surrounding area.
- ◆ Seborrheic dermatitis:
- apply to the affected area once or twice daily depending on the severity of infection.
- ◆ Treatment should be continued for a sufficient period, at least until a few days after disappearance of all symptoms. The diagnosis should be reconsidered if no clinical improvement is noted after 4 weeks of treatment. General measures in regard to hygiene should be observed to control sources of infection or reinfection.
- ◆ The usual duration of treatment is : tinea versicolor (white spots) 2-3 weeks, yeast infections 2-3 weeks, jock itch 2-4 weeks, ringworm of the body 3-4 weeks, athlete's foot 4-6 weeks, seborrheic dermatitis 2-4 weeks.
- ◆ Maintenance therapy is applied once or twice weekly in seborrheic dermatitis.

Side Effects/Adverse Reactions : A few instances of irritation and burning sensation have been observed during treatment with Ketoconazole cream.

Contraindications : Known sensitivity to ketoconazole or the excipients.

Precautions/Warnings : Topical use of ketoconazole is generally well-tolerated.

Carcinogenicity :

A long term feeding study in Swiss albino mice and Wistar rats has shown no evidence in carcinogenicity.

Mutagenicity :

The dominant lethal mutation test in male and female mice given oral ketoconazole, has shown no mutation at any stage of germ cell development.

Breast-feeding :

No systemic absorption was detected following a single application to the chest, back, and arms of healthy volunteers. Therefore, topical ketoconazole is unlikely to be excreted in breast milk in significant amounts, or to cause adverse effects in the nursing infant. Not for ophthalmic use and use in nursing mothers and paediatric. Do not use on children under 2 years of age except under the advice and supervision of a doctor. Avoid contact with eyes. Consult the doctor if no improvement in 4 weeks (for athletes' foot) & 2 weeks (for jock itch).

Overdosage and Treatment : Exaggerated topical application may lead to erythema, oedema and a burning sensation, which will disappear upon discontinuation of the treatment. In the event of accidental ingestion, supportive measures should be carried out.

Storage : Store below 30°C. For External Use only.

Presentation : White, smooth, homogeneous and water-miscible cream in 15g collapsible aluminium tube.

KEEP ALL MEDICINES OUT OF REACH OF CHILDREN
JAUHI DARI KANAK-KANAK

Shelf-life : The expiry date is indicated on the packaging.

Revision Date : 08-Dec-2017



Xepa-Soul Pattinson (Malaysia) Sdn Bhd Company No.8504-M 1-5 Cheng Industrial Estate, 75250 Melaka, Malaysia.

