

HOVID GENTAMICIN CREAM 0.1%

VIGEN05-var (MY)

DESCRIPTION

White, opaque cream.

COMPOSITION

Gentamicin (as Sulphate) 0.1% w/w.

PHARMACODYNAMICS

Gentamicin is an aminoglycosides and bactericidal. It is actively transported across the bacterial cell membrane, binds to a specific receptor protein on the 30 S subunit of bacterial ribosomes and interferes with an initiation complex between mRNA (messenger RNA) and the 30 S subunit, inhibiting protein synthesis. DNA may be misread, thus producing nonfunctional proteins; polyribosomes are split apart and are unable to synthesize protein.

PHARMACOKINETICS

Although not absorbed through intact skin, topical gentamicin is readily absorbed from larger denuded, burned or granulating areas. Greater and more rapid absorption occurs with gentamicin cream than with the ointment. When gentamicin is applied to large areas of denuded (e.g. burned) body surface, plasma concentrations can reach 1mg/ml. Plasma elimination half-life for gentamicin is 2 to 3 hours and may be considerably longer in neonates and patients with renal impairment. Gentamicin is excreted in urine.

INDICATIONS

Treatment of impetigo erythema and other localised primarily bacterial skin infection with a Gram-negative component and also for secondary bacterial infections complicating other pre-existing dermatoses.

CONTRAINDICATIONS

Gentamicin is contraindicated in patients with a known history of allergy to it and probably to other aminoglycosides.

As this preparation contains propylene glycol, it should be avoided in patients who are hypersensitive to the propylene glycol.

WARNINGS AND PRECAUTIONS

The overgrowth of nonsusceptible organisms, including fungi occasionally occurs with the use of topical antibiotics. If this occurs, or irritation, sensitization or superinfection develops, treatment should be discontinued and consult physician.

When gentamicin is applied topically to large denuded areas of skin, the small risk of ototoxicity from systemic absorption should be considered, especially in the elderly and in those with impaired renal function.

Use of topical gentamicin preparation in closed hospital settings is actively discouraged

PREGNANCY AND LACTATION

Consideration should be given to the possibility of foetal ototoxicity when gentamicin is applied topically to large denuded areas of skin.

MAIN SIDE/ADVERSE EFFECTS

The following side-effects resulting in hypersensitivity reactions may occur:

- itching
- redness
- swelling
- other sign of irritation not present before therapy

OVERDOSE AND TREATMENT**Clinical features**

Redness and itching of skin

Treatment

Discontinuation of therapy will normally bring about relief to the symptoms

DOSAGE AND ADMINISTRATION**Adults**

Topical, to the skin, 3 or 4 times a day.

Paediatric

Children over 1 year of age - see adult dose.

Note: The information given here is limited. For further information, consult your doctor or pharmacist.

Storage:

Store below 30°C.

Presentation/Packing:

Cream 0.1% x 15 g

Product Registration Holder/

Manufactured by: HOVID Bhd.
121, Jalan Tunku Abdul Rahman
(Jalan Kuala Kangsar),
30010 Ipoh, Perak, Malaysia.

Revision date: June 2021