

YSP NYSTATIN

CAPSULE 500,000units

Nystatin is a polyene antifungal agent derived from cultures of a strain of *Streptomyces noursei*. It is used for the prophylaxis and treatment of candidiasis of the skin and mucous membranes. Nystatin is less affected by changes in pH than other antifungal drugs.

Ingredient(s):

Each capsule contains:

Nystatin 500,000 units

Pharmacodynamics:

1. Nystatin exerts its antifungal effects by binding to sterols in the fungal cell membrane to produce a change in membrane permeability which allows loss of essential cellular constituents.
2. Nystatin has both fungistatic and fungicidal effects against a wide variety of yeast and yeast-like fungi. Its main action is against *Candida* spp. It is also effective against *Aspergillus* spp., *Coccidioides immitis*, *Cryptococcus neoformans*, *Histoplasma capsulatum*, *Blastomyces dermatitidis*, and other yeasts and fungi.

Pharmacokinetic:

1. Nystatin is poorly absorbed from the GI tract, and detectable blood concentrations are not obtained after usual doses. Following oral administration, Nystatin is excreted almost entirely in feces as unchanged drug. The drug is not absorbed from intact skin or mucous membranes.

Indication(s):

For the prevention and treatment of intestinal candidiasis.

Route of Administration:

Oral

Dosage and Administration:

The usual therapeutic dosage is one to two capsules (500,000 to 1,000,000 units Nystatin) three times daily. Treatment should generally be continued for at least 48 hours after clinical cure to prevent relapse. To be used on physician's prescription.

Contraindication(s):

This preparation is contraindicated in patients with a history of hypersensitivity to Nystatin.

Precaution(s) / Warning(s):

No adverse effects or complications have been attributed to Nystatin in infants born to women treated with Nystatin.

Side Effect (s) / Adverse Reaction(s):

Nystatin is virtually nontoxic and nonsensitizing and is well tolerated by all age groups, including debilitated infants, even on prolonged administration. Large oral doses have occasionally produced diarrhea, gastrointestinal distress, nausea, and vomiting. Other side effects reported are oral irritation and sensitization, rash (including urticaria) and rarely Steven-Johnson Syndrome.

Symptoms and Treatment for Overdosage, and Antidote(s):

Following oral administration, Nystatin is only sparingly absorbed, with no detectable blood levels when given in the recommended doses. Therefore no risk of overdosage exists.

Shelf-Life:

3 years from the date of manufacture.

Storage Condition(s):

Keep in a tight container. Store at temperature below 30°C. Protect from light and moisture.

Product Description and Packing(s):

A deep brownish red cap and pale pink body hard gelatin capsule, containing yellow powder.

Plastic container of 200's and 1000's (for export only).

Plastic container of 100's and 180's.



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679