

DYNA AMPICILLIN 125MG/5ML

MAL19985980AZ

DESCRIPTION:

Dry Syrup

Colour : White (Yellow on adding water)

Flavour : Strawberry

CONTENT:

Each 5 ml of the reconstituted suspension contains:

Ampicillin Trihydrate equivalent to Ampicillin 125 mg

Preservative: Sodium Benzoate 0.1% w/v

PHARMACODYNAMICS:

Ampicillin is a bactericidal antibiotic which inhibits bacterial cell wall synthesis, probably by acylation of membrane-bound transpeptidase enzymes. This prevents cross-linkage of peptidoglycan chains which is necessary for bacterial cell wall strength and rigidity. Also, cell division and growth are inhibited and lysis of susceptible bacteria frequently occurs. Rapidly dividing bacteria are most susceptible to penicillin group of antibiotics including Ampicillin.

PHARMACOKINETICS:

Ampicillin is acid stable and 30 to 60% of an oral dose is absorbed. Peak serum concentrations of 0.8 to 4 mcg/ml are attained 1 to 2 hours after oral doses of 250 to 500 mg of anhydrate or trihydrate. The serum half-life is 1 to 2 hours which is increased in subjects with renal function impairment. Ampicillin enters most body cavities, tissues and fluids except the cerebrospinal fluid unless the meninges are inflamed, in which case large amounts may be detectable in this fluid; Ampicillin crosses the placenta and is excreted in the milk; volume of distribution about 20 L which in subjects with cirrhosis may reach about 60 L. About 20% of Ampicillin is bound to plasma proteins. 30% to 50% of an oral dose is excreted in the urine in 6 to 8 hours; up to 70% is excreted in 24 hours; urinary excretion is decreased after the administration of probenecid; Ampicillin is excreted in the bile in concentrations up to 4 mcg/ml after an oral dose of 250 mg; biliary excretion is increased in subjects with cirrhosis.

INDICATIONS:

Infections caused by sensitive gram-positive organisms, including *Streptococcus pneumoniae*, and other streptococci and *Listeria monocytogenes*. Gram-negative microorganism including some of strains of *Bordetella pertussis*, *Haemophilus influenzae* and some Enterobacteriaceae such as *Escherichia coli*, *Proteus mirabilis*, *Salmonella* and *Shigella* spp.

Gram-negative cocci such as *Branhamella catarrhalis*, *Neisseria gonorrhoeae*, and *N. Meningitidis*.

ENT infections: otitis media, sinusitis, tonsillitis, pharyngitis, laryngitis, epiglottitis, and acute bronchitis

Lower respiratory tract infections: chronic bronchitis, pneumonia, pleuritis.

Urinary tract infections: acute and recurrent cystitis, acute and chronic pyelonephritis, asymptomatic bacteriuria, prostatitis.

Skin and soft tissue infections: Erysipelas, impetigo, pyoderma.

Gonorrhoea and enteric infections.

RECOMMENDED DOSE:

Respiratory-tract or skin infections:

Children 2 to 10 years : 125 mg to 250 mg every 6 hours.

Children < 2 years : 62.5 mg to 125 mg every 6 hours.

Gastro-intestinal or urinary-tract infections:

Children 2 to 10 years : 125 mg to 250 mg every 6 hours.

Children < 2 years : 62.5 mg to 125 mg every 6 hours.

For gonorrhoea:

3 g as a single oral dose may be administered with 1 g probenecid.

The dose should be reduced in renal failure.

DIRECTION FOR MIXING:

Shake to loosen powder.

Add approximately half the final volume (30 ml or 50 ml) of freshly boiled and cooled water and shake gently.

Add more water until the final volume (60 ml or 100 ml) mark is reached and shake well.

Keep refrigerated (2-8°C) after reconstitution. To be used within 7 days of mixing. **Complete** the prescribed course.

ROUTE OF ADMINISTRATION:

FOR ORAL USE ONLY

CONTRAINDICATIONS:

Not to be used in patients with known hypersensitivity to penicillin.

WARNINGS AND PRECAUTIONS:

Patients hypersensitive to other penicillins or cephalosporins or penicillamine may be hypersensitive to Ampicillin. Reduced dose may be required in patients with impaired renal function.

Serious and occasionally fatal hypersensitivity reactions (including anaphylactoid and severe cutaneous adverse reactions) have been reported in patients on penicillin therapy with beta-lactams. Before initiating therapy with Dyna Ampicillin 125mg/5ml, careful inquiry should be made concerning previous hypersensitivity reactions to penicillins, cephalosporins, carbapenems or other beta-lactam agents. If an allergic reaction occurs, Dyna Ampicillin 125mg/5ml must be discontinued immediately and appropriate alternative therapy instituted.

DRUG INTERACTIONS:

Probenecid prolongs blood levels of Ampicillin. Antacids and other alkalisng agents can impair absorption of Ampicillin. May reduce effectiveness of oral contraceptives. Concurrent administration with allopurinol increases incidence of rashes. It decreases the efficacy of oestrogen containing oral contraceptives. It may also affect the absorption of other drugs due to its effect on the gastro-intestinal flora.

PREGNANCY AND LACTATION:

Risk-benefit must be considered when given to pregnant women and during breast-feeding.

SIDE EFFECTS:

Hypersensitivity or allergic reactions i.e. skin rashes either urticarial or maculopapular, itching or wheezing.

GIT irritation i.e. abdominal or stomach cramps, pain, bloating, nausea, vomiting, diarrhoea and pseudomembranous colitis.

Fever, increased thirst, unusual tiredness or weakness.

Supra-infection with *Pseudomonas* & *Candida* with prolonged use.

Skin and subcutaneous tissue disorders:

Frequency 'very rare': Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS).

SYMPTOMS AND TREATMENT OF OVERDOSE:

Problems of overdose are unlikely to occur. If encountered, they should be treated symptomatically. Ampicillin may be removed from circulation by haemodialysis.

PACKING/PACK SIZE(S):

Plastic bottles of 60 ml and 100 ml.

JAUHI UBAT DARIPADA KANAK-KANAK

KEEP OUT OF REACH OF CHILDREN

SHAKE WELL BEFORE USE

MANUFACTURER/PRODUCT REGISTRATION HOLDER:

DYNAPHARM (M) SDN. BHD. 198001011897 (65683-V)

2497, Mk. 1, Lorong Perusahaan Baru 5,

Kawasan Perusahaan Perai 3,

13600 Perai, Pulau Pinang, MALAYSIA

P11PL A001-05

Latest Revision: Oct 2023