

CASPRIN ENTERIC MICROENCAPSULATED CAPSULE

Ingredient(s):

Each vegetable capsule contains:

Aspirin 100mg

Pharmacodynamic:

1. Aspirin is an analgesics, anti-inflammatory, antipyretic and an inhibitor of platelet aggregation. It acetylates the enzymes cyclo-oxygenase thereby inhibiting the synthesis of prostaglandins, thromboxane in the circulation platelets, and prostacyclin in the systemic vascular endothelium. Thromboxane is a vasoconstrictor and it induces platelet aggregation, while prostacyclin is a vasodilator and inhibitor of platelet aggregation.
2. The antiplatelet aggregating effect of Casprin is postulated to retard or reduce the development of thromboembolic disorders. The low dose, enteric coated, and relatively slow rate of release of Aspirin is observed to selectively and irreversibly inhibit thromboembolic synthesis. Aspirin inhibits thromboxane synthesis in the portal circulation and is presystemically metabolised to salicylate with negligible amount reaching the systemic circulation to inhibit prostacyclin synthesis. Salicylate is effectively inactive at the blood concentrations achieved clinically and prostacyclin is less sensitive than thromboxane to Aspirin.

Pharmacokinetics:

1. In order to eliminate the gastrointestinal adverse effects of Aspirin, an enteric coated low-dose Aspirin (as pellets) was developed by Yung Shin Pharmaceuticals. The advantages of enteric-microencapsulated capsules include ready distribution over a large area, thus minimizing the risk of local damage caused by the dumping effect of enteric coated tablets. Furthermore, Casprin Enteric-Microencapsulated Capsules are also less dependent on gastric transient time. They may attain more constant plasma level, achieve slow-release effect, give higher accuracy in reproducibility between doses, and provide less decreases in bioavailability.
2. Following oral administration, absorption of Aspirin occurs in the intestine. Some Aspirin is hydrolysed to salicylate in the gut wall. Aspirin is bound to plasma protein and is widely distributed. Plasma Aspirin concentrations decline rapidly (half life 15 to 20 minutes) as plasma salicylates concentrations increases.
3. Aspirin is rapidly converted by esterases present in plasma and many tissues, especially the liver, to salicylic acid which itself has some antipyretic, analgesic and anti-inflammatory actions, but which has little effects on platelets. Salicylic acid is metabolised in the liver to the glycine conjugate salicyluric acid. Other metabolites include salicyl phenolic glucuronide, salicyl acyl glucuronide and gentisic acid. Excretion of salicylic acid in urine is pH dependent; approximately 80% appears unchanged in urine at pH 8, but only around 10% at pH 4.
4. Since enteric coated pellets of Aspirin can reduce gastrointestinal side effects that may occur with conventional or other enteric coated tablet preparations, it is an effective antiplatelet treatment of choice for long term treatment and prevention of certain forms of ischemic attacks.

Indication(s):

In the treatment of acute myocardial infarction (heart attack), for prevention of heart attack in certain high-risk group patients and for prevention of stroke after transient ischemic attack (TIA).

Dosage and Administration:

One capsule daily to be taken with a glass of water.

Contraindication(s):

Severe hepatic and renal disease, haemophilia or other bleeding disorders, erosive gastritis or peptic ulcers. Patients who are allergic to aspirin.

Side Effect(s) / Adverse Reaction(s):

The main adverse reactions associated with Aspirin therapy include gastrointestinal distress, nausea, vomiting, erosion of the gastric mucosa, ulceration and occult blood loss.

Precaution(s) / Warning(s):

1. Because of its relatively low dosage and enteric coating dosage form, the risk of Casprin in causing GI ulceration is greatly reduced. However, care should be taken in administering Casprin to patients with prior history of serious GI events.
2. Concurrent use of Casprin with alcoholic beverages should be avoided in order to avoid GI toxicity.
3. Care should be taken when Aspirin is administered to asthmatics patients.
4. Caution is necessary when renal or hepatic function is impaired.
5. The use of Aspirin in children under the age of 16 years is not recommended because of the risk of Reye's syndrome.

Pregnancy and Lactation:

1. Pregnancy: May cause premature closure of the fetal ductus arteriosus; avoid during 3rd trimester. May cause excessive blood loss at delivery; avoid 1 week prior to and during labor/delivery.

2. Lactation: Not for use in nursing.

Drug Interaction(s):

- 1. Concurrent administration of Aspirin and dipyridamole may result in an increase in peak plasma-salicylate concentration and area under the curve.
- 2. Serum-salicylate concentrations may be reduced by concurrent administration of corticosteroids.
- 3. Aspirin may enhance the activity of coumarin anticoagulant, sulphonylurea, hypoglycaemic agent, methotrexate, phenytoin and valproic acid.

Symptoms and Treatment for Overdosage and Antidote(s):

Mild cases may produce nausea, vomiting, sweating, thirst and tachycardia. Severe overdosage may show fever and CNS disturbances such as convulsions, hallucinations, coma and respiratory failure. Management includes gastric lavage and emesis for mild cases; in severe cases, barbiturate administrations or dialysis may be required.

Shelf-Life:

2 years from the date of manufacture.

Storage Condition(s):

Store at temperature below 30°C. Protect from light and moisture.

Product Description and Packing(s):

A size #4 orange transparent cap and colorless transparent body vegetable capsule, containing white to off-white pellets, both cap and body impressed with the mark "AECp". Blister packing of 10's x 10, 10's x 3, 30's, 30's x 2 and 30's x 50.

CASPRIN MIKROPENGKAPSULAN ENTERIK KAPSUL

Bahan Aktif:

Setiap kapsul sayuran mengandungi:

Aspirin 100mg

Indikasi:

Digunakan untuk serangan jantung dan pencegahan serangan jantung bagi pesakit-pesakit dalam kumpulan yang berisiko tinggi. Ia juga digunakan untuk pencegahan strok selepas serangan iskemia sementara.

Dos:

1 kapsul sehari, diambil bersama satu gelas air.

CASPRIN ENTERIC MICROENCAPSULATED CAPSULE

成份：

每顆素食膠囊含有：

Aspirin 100mg

適應症：

預防及治療短暫缺血性發作（小中風）及其他血栓性栓塞疾患。

用法・用量：

每日服用一粒，與溫水服用。



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679