



## DEZOR CREAM

### NAME AND STRENGTH OF ACTIVE SUBSTANCE:

Ketoconazole 2% w/w.

### PRODUCT DESCRIPTION:

A white cream.

Other ingredients are Chlorocresol, Butylated Hydroxytoluene, Hydrochloric Acid 37%, Dibasic Sodium Phosphate (Disodium Hydrogen Phosphate Dihydrate), Polyethylene Glycol 400, White Soft Paraffin, Cetomacrogol Emulsifying Wax, Liquid Paraffin and Purified Water.

**PHARMACODYNAMICS :** Ketoconazole is an imidazole antifungal agent which interferes with ergosterol synthesis and therefore alters the permeability of the cell membrane of sensitive fungi.

**PHARMACOKINETICS :** Ketoconazole is a broad-spectrum synthetic antifungal agent which inhibits the growth of the following common dermatophytes and yeasts by altering the permeability of the cell membrane : *Dermatophytes-Trichophyton rubrum, T. mentagrophytes, T. tonsurans, Microsporum canis, M. audouini, M. gypseum and Epidermophyton floccosum; Yeasts-Candida albicans and C. tropicalis* and the organism responsible for tinea versicolor, *Malassezia furfur (Pityrosporum orbiculare)*. Development of resistance to Ketoconazole has not been established. *In vitro* studies suggest that Ketoconazole impairs the synthesis of ergosterol which is a vital component of fungal cell membranes. The mechanism of therapeutic effect of Ketoconazole in seborrheic dermatitis has not been established.

**INDICATIONS :** Topical treatment of tinea corporis, tinea cruris, tinea pedis, pityriasis versicolor and cutaneous candidiasis.

**CONTRAINDICATIONS :** Ketoconazole cream is contra-indicated in individuals who have shown hypersensitivity to any of its ingredients. Ketoconazole cream is not for ophthalmic use.

**DRUG INTERACTIONS :** Concomitant administration of Ketoconazole with rifampicin results in decreased antifungal activity. Concurrent use of Ketoconazole and Phenytoin may alter serum concentrations of both drugs.

**ADVERSE EFFECTS :** Signs of skin irritation-burning sensation, pruritus, dermatitis were observed in 3% of Ketoconazole treated patients and in 2% of placebo controls during clinical trials.

**WARNINGS AND PRECAUTIONS :** If irritation or sensitivity occurs, treatment should be discontinued. Do not use on children under 2 years of age except under the advice and supervision of a doctor. Avoid contact with eyes and mucous membranes (for external use only). Consult doctor if there is no improvement in 4 weeks (for athlete's foot) or 2 weeks (for jock itch). Overgrowth of non susceptible organisms may occur with prolonged use of any topical antimicrobial agent including Ketoconazole.

**Use in pregnancy and lactation:** Animal studies and studies in human volunteers have failed to demonstrate systemic absorption of topically applied Ketoconazole cream; however due to the teratogenic nature of the active ingredient, Ketoconazole cream should only be used in pregnant or lactating woman if in the judgement of the treating physician, the expected benefits outweigh the potential risk.

**ROUTE OF ADMINISTRATION :** Topical.

**RECOMMENDED DOSAGE :** Apply once daily. Duration of treatment: Pityriasis versicolor, yeast infection, tinea cruris, Tinea corporis 2 weeks and Tinea pedis 4 - 6 weeks.

### SYMPTOMS AND TREATMENT FOR OVERDOSE :

No known information available concerning overdosage in human.

**PACKING :** 5g and 15g collapsible aluminium tubes.  
450g in polyethylene jars (for export only).  
Not all presentations may be available locally.

**STORAGE :** Store below 30°C. For external use only. Keep out of reach of children.  
Recommended shelf-life: 3 years.

### MANUFACTURED BY :

**HOE Pharmaceuticals Sdn. Bhd.**

Lot 10, Jalan Sultan Mohamed 6, Bandar Sultan Suleiman,  
42000 Port Klang, Selangor, MALAYSIA.

Revision date :August 2022