

pharmaniaga®



Ziconal Cream

Ketoconazole BP 2%

DESCRIPTION

A smooth white cream.

COMPOSITION

Ketoconazole 2% w/w.

Ketoconazole cream also contains chlorocresol 0.1% w/w as preservative, and propylene glycol as diluent.

ACTIONS

Ketoconazole, a synthetic imidazole dioxolane derivative, has a potent antimycotic activity against dermatophytes such as *Trichophyton* spp., *Epidermophyton floccosum* and *Microsporum* spp. and against yeasts, including *Malassezia* spp. and *Candida* spp. The effect on *Malassezia* spp. is particularly pronounced.

Plasma concentrations of ketoconazole were not detectable after topical administration of ketoconazole 2% cream in adults on the skin.

INDICATIONS

Ketoconazole cream is indicated for the topical treatment of tinea corporis, tinea cruris and tinea pedis caused by *Trichophyton rubrum*, *T. mentagrophytes* and

Epidermophyton floccosum; in the treatment of tinea versicolour caused by *Malassezia furfur*, in the treatment of cutaneous candidiasis caused by *Candida* spp. and the treatment of seborrheic dermatitis.

CONTRAINDICATION

Ketoconazole cream is contraindicated in patients hypersensitivity to ketoconazole or propylene glycol.

ADVERSE REACTIONS

Side-effects include itching, stinging or irritation.

PRECAUTIONS

If a reaction suggesting sensitivity or chemical irritation should occur, use of the medication should be discontinued. Hepatitis (1 : 10,000 reported incidence) and, at high doses, lowered testosterone and ACTH induced corticosteroid serum levels have been seen with orally administered ketoconazole; these effects have not been seen with topical ketoconazole.

Pregnancy

Ketoconazole crosses the placenta.

Ketoconazole should be used during pregnancy only if the potential benefit justifies the potential risk to the foetus.

Breast-feeding

It is not known whether ketoconazole cream administered topically could result in sufficient systemic absorption to produce detectable quantities in breast milk.

However, no systemic absorption was detected following a single application to the chest, back and arms. Therefore, topical ketoconazole is unlikely to be distributed into breast milk in significant amounts as to cause adverse effects in the nursing infant.

DRUG INTERACTIONS

None known.

Effects on Ability to Drive and Use Machine

Ketoconazole cream has no influence on the ability to drive and use machines.

RECOMMENDED DOSAGE

Cutaneous candidiasis, tinea corporis, tinea pedis and tinea (pityriasis) versicolour. It is recommended that ketoconazole cream be applied once daily to cover the

affected and immediate surrounding area. Clinical improvement may be seen fairly soon after treatment is begun; however, candidal infections and tinea cruris and corporis should be treated for two weeks in order to reduce the possibility of recurrence. Patients with tinea versicolour usually require two weeks of treatment.

Seborrheic dermatitis: Ketoconazole cream should be applied to the affected area twice daily for four weeks or until clinical clearing.

If a patient shows no clinical improvement after the treatment period, the diagnosis should be redetermined.

OVERDOSAGE

Ketoconazole cream is not absorbed systemically in detectable levels.

ROUTE OF ADMINISTRATION

Topical.

STORAGE CONDITIONS

Store below 30°C.
Protect from light.
Keep container tightly closed.

SHELF LIFE

Product should not be used beyond the expiry date imprinted on the product packaging.

PRESENTATION

Tubes of 15g.

Date of revision: 28 October 2024

PRODUCT REGISTRATION HOLDER:

Pharmaniaga Manufacturing Berhad (198001006232)
No. 11A, Jalan P/1, Kawasan Perusahaan Bangi,
43650 Bandar Baru Bangi,
Selangor Darul Ehsan, Malaysia.

MANUFACTURER:

Idaman Pharma Manufacturing Sdn Bhd (200401023395)
Lot 24 & 25, Jalan Perusahaan Lapan,
Bakar Arang Industrial Estate,
08000 Sungai Petani, Kedah, Malaysia.

Lt-404.00

115mm

150mm