

Kezoral

2% cream

DESCRIPTION

Kezoral Cream: White colour, homogenous cream containing Ketoconazole 2%.
Preservatives : Methylparaben 0.25%
Propylparaben 0.048%

PHARMACODYNAMICS

Ketoconazole is fungistatic, and may be fungicidal, depending on concentration. Ketoconazole interferes with cytochrome P-450 activity, which is necessary for the demethylation of 14- α -methylsterols to ergosterol. Ergosterol, the principal sterol in the fungal cell membrane, becomes depleted. This damages the cell membrane, producing alterations in membrane functions and permeability. In *Candida albicans*, ketoconazole inhibits transformation of blastospores into invasive mycelial form.

INDICATIONS

Indicated as a primary agent in the topical treatment of dermatophyte infections of the skin: Tinea corporis (ringworm of the body) and tinea cruris (ringworm of the groin; jock itch) caused by *Trichophyton rubrum*, *T. mentagrophytes* and *Epidermophyton floccosum* (*Acrotheium floccosum*), tinea pedis (athlete's foot), as well as in the treatment of pityriasis versicolor (tinea versicolor; "sun fungus") caused by *Malassezia furfur* (*Pityrosporum orbiculare*) and cutaneous candidiasis caused by *Candida* species. Kezoral Cream can be used in the treatment of seborrhoeic dermatitis.

PHARMACOKINETICS

No systemic absorption detected at a sensitivity level of 5 nanogram per ml in the blood over a 72-hour period following a single topical application to the chest, back and arms of normal volunteers.

DOSAGE AND ADMINISTRATION

Kezoral cream should be applied in sufficient quantity to cover affected and surrounding areas, the cream gently rubbed in to enhance its effect. The patient must be counselled in regards to compliance with the full course of therapy and that some fungal infections may require prolonged therapy.

Type Of Infection	Frequency Of Application	Length Of Therapy
Cutaneous candidiasis, Tinea corporis, Tinea cruris, Pityriasis versicolor	Once a day	At least 2 weeks
Tinea pedis	Once a day	At least 6 weeks
Seborrhoeic dermatitis	Once to twice a day	At least 4 weeks or until clinical clearing

TOXICOLOGY

Cross-sensitivity and/or related problems

Person sensitive to miconazole or other imidazoles may be sensitive to ketoconazole also.

Carcinogenicity

A long-term feeding studies in Swiss albino mice and Wistar rats has shown no evidence of carcinogenicity.

Mutagenicity

The dominant lethal mutation tests in male and female mice, given single oral doses of ketoconazole as high as 80mg per kg of body weight (mg/kg), has shown no mutations at any stage of germ cell development. The Ames Salmonella microsomal activator assay has also shown negative result.

Pediatrics

No information is available on the relationship of age to the effects of this medicine in pediatric patients. Safety and efficacy have not been established.

Geriatrics

No information is available on the relationship of age to the effects of this medicine in geriatric patients.

Pregnancy/Reproduction

Pregnancy-Ketoconazole crosses the placenta. Adequate and well-controlled studies in humans may not been done. Studies in rats, given oral doses of 80 mg/kg per day (10 times the maximum recommended human dose [MRHD]), have shown ketoconazole to be teratogenic, causing syndactyly and oligodactyly. FDA Pregnancy Category C.

Breast-feeding

It is not know whether ketoconazole, applied topically on a regular basis, is absorbed systemically in sufficient amounts to be distributed into breast milk in detectable quantities. However, no systemic absorption was detected following a single application to the chest, back and arms of healthy volunteers. Therefore, topical ketoconazole is unlikely to be distributed into breast milk in significant amounts or to cause adverse effects in the nursing infant.

CONTRAINDICATIONS

Hypersensitivity to propylene glycol and ketoconazole.

PRECAUTIONS

Kezoral cream is intended for application to unbroken skin only and is not for ophthalmic use.

SIDE EFFECT

Infrequent reports of skin irritation (itching and stinging) have been reported. In rare cases, prolonged use may result in irritation of the skin which may require discontinuation of treatment.

OVERDOSAGE AND TREATMENT

Exaggerated use can bring on local irritation and erythema. Discontinuation of treatment remedies this effect.

PRESENTATION:

Cream: In packs of 15g, 30g.

All pack size(s) stated may not be available locally

Store in a dry place below 30°C
FOR EXTERNAL USE ONLY
Keep container tightly closed
Keep out of reach of children
Jauhi daripada kanak-kanak
Shelf life: Please refer to outer package.

Product Registration Holder:

Duopharma Manufacturing (Bangi) Sdn Bhd.

Lot No. 2, 4, 6, 8 & 10, Jalan P/7, Section 13,
Bangi Industrial Estate, 43650 Bandar Baru Bangi,
Selangor, Malaysia.

Manufacturer :

Duopharma Manufacturing (Bangi) Sdn Bhd.

Lot No. 2 & 4, Jalan P/7, Section 13,
Bangi Industrial Estate,
43650 Bandar Baru Bangi,
Selangor, Malaysia.



1500008660 N1
Revision Date: May 2019

IKEZORAL CI

IKEZORAL CI