

PACKAGE INSERT

SUNAZOL CREAM

Composition: Ketoconazole _____ 2%/w/w

Product Description: White and smooth cream.

Pharmacodynamics:

Ketoconazole is a broad spectrum synthetic antifungal agent which inhibits the in vitro growth of the following common dermatophytes and yeasts by altering the permeability of the cell membrane. In vitro studies suggest that ketoconazole impairs the synthesis of ergosterol, which is a vital component of fungal cell membranes. It is postulated that the therapeutic effect of ketoconazole in seborrheic dermatitis is due to the reduction of *M. ovale*, but this has not been proven.

Pharmacokinetics:

When ketoconazole 2% cream was applied dermally to intact or abraded skin of Beagle dogs for 28 consecutive days at a dose of 80 mg, there were no detectable plasma levels using an assay method having a lower detection limit of 2 ng/ml. After a single topical application to the chest, back and arms of normal volunteers, systemic absorption of ketoconazole was not detected at the 5ng/ml level in blood over a 72-hour period.

Indication:

It is indicated for the topical treatment of tinea corporis, tinea cruris, tinea manus, tinea pedis, cutaneous candidiasis and pityriasis versicolor. Also indicated for the treatment of seborrheic dermatitis, a skin condition related with the presence of *Pityrosporum ovale*.

Recommended Dosage:

For treatment of tinea corporis, tinea cruris, tinea manus, tinea pedis, cutaneous candidiasis and pityriasis versicolor, it is recommended to apply ketoconazole 2% cream to the affected and immediate surrounding area once daily. The usual duration for the treatment of tinea corporis is 3 - 4 weeks, tinea cruris is 2 - 4 weeks, tinea pedis is 4 - 6 weeks and yeast infection is 2 - 3 weeks. For treatment of seborrheic dermatitis, it is recommended to apply to the affected area once or twice daily depending on the severity of the infection. The usual duration of treatment is 2 - 4 weeks. Ketoconazole cream is applied once or twice weekly for maintenance treatment of seborrheic dermatitis. The cream should be applied for a sufficient period, at least until a few days after disappearance of all symptoms. If there is no clinical improvement after a month of treatment, further medical consultation should be done.

Route of Administration: Topical

Contraindications:

Ketoconazole is contraindicated in individuals who have shown hypersensitivity to it and propylene glycol.

Warnings and Precautions:

If a reaction suggesting sensitivity or chemical irritation should occur, use of ketoconazole should be discontinued. After a prolonged treatment with topical corticosteroids, irritation may occur if ketoconazole is applied immediately. Hence, it is advisable to continue the treatment of the topical corticosteroid in the morning and to apply ketoconazole cream in the evening and to subsequently and gradually withdraw the steroid treatment over a period of 2 to 3 weeks. If a potent steroid is applied, replace it by a mild steroid and withdraw the treatment over a period of 2 to 3 weeks also. Ketoconazole cream must not be applied on the eyes.

Interactions with Other Medicaments:

No known drug interaction since ketoconazole cream applied topically is not absorbed.

Pregnancy and Lactation:

There are no adequate and well-controlled studies and ketoconazole should be used only if the potential benefit justifies the potential risk.

Side Effects:

Reported side effects are mainly of irritation, pruritus and stinging.

Symptoms and Treatment of Overdose:

Exaggerated use can bring on local irritation and erythema. Discontinuation of treatment remedies this effect.

Storage Conditions: Store at temperature 30°C. Protect from light.

Pack Sizes : A tube of 15g and 30g.

Pack Sizes (export only) : A jar of 400g.

Shelf-life: 2 years.

FURTHER INFORMATION CONCERNING THIS DRUG CAN BE OBTAINED FROM YOUR FAMILY PHYSICIAN / LOCAL GENERAL PRACTITIONER / PHARMACIST.

Manufacturer & Product Registration Holder:

SUNWARD PHARMACEUTICAL SDN. BHD.

No. 3, 11 & 17, Jalan Kempas 4,

Taman Petrodustrian Tampoi Indah,

81200 Johor Bahru, Johor, Malaysia.

EL221A-R9

Revision Date : 24/11/2005