

	Client		Color : BLACK 	Finishing :
	Product	Dirlip BIFEN 100mg MY		
	Dimension	110 x 260mm		
	Code	40985-03		
	Material			
	Date			

BIFEN SUSPENSION 100 mg/5 ml

DESCRIPTION:

Orange suspension with orange flavour and sweet taste.

COMPOSITION:

Ibuprofen BP 100 mg / 5ml

PHARMACOLOGY:

Ibuprofen is a non-steroidal anti-inflammatory that possesses analgesic and anti-pyretic activities. It inhibits the activity of the enzyme cyclo-oxygenase, resulting in decreased formation of precursors of prostaglandins and thromboxanes from arachidonic acid.

PHARMACOKINETIC:

Approximately 80% of an oral dose of ibuprofen is absorbed from the GI tract. About 90-99% of a dose is bound to plasma proteins. The plasma half-life of the drug has been reported to be 2-4 hours. Excretion of ibuprofen is essentially completed within 24 hours.

INDICATION:

Children : Symptomatic relief of pain and fever associated with colds, influenza, toothache, bronchitis and other bacterial and viral infections.

Adults : Management of mild to moderate pain in conditions such as dysmenorrhoea, migraine, postoperative pain, ankylosing spondylitis, osteoarthritis and rheumatoid arthritis and in other musculoskeletal and joint disorders.

ADVERSE EFFECT:

Gastrointestinal disturbances, peptic ulceration, gastrointestinal bleeding. Others include headache, dizziness, nervousness, skin rash, pruritus, tinnitus, edema, depression, drowsiness, insomnia, blurred vision, hypersensitivity, impairment of renal function and thrombocytopenia.

PRECAUTIONS:

Ibuprofen should be given with care to patient with asthma or bronchospasm, bleeding disorders, cardiovascular disease, a history of peptic ulceration and in liver and renal failure. Care is required in those who are receiving coumarin anticoagulants. Ibuprofen should be discontinued

in patients who experience blurred or diminished vision or changes in colour vision. Patients with collagen disease may be at increased risk of developing aseptic meningitis.

WARNING:

RISK OF ULCERATION, BLEEDING AND PERFORATION WITH NSAID

Serious GI toxicity such as bleeding, ulceration and perforation can occur at any time, with or without warning symptoms in patients treated with NSAID therapy. Although minor GI problems (eg. Dyspepsia) are common, usually developing early in therapy, prescribers should remain alert for ulceration and bleeding in patients treated with NSAIDs even in the absence of previous GI tract symptoms. In patients observed in clinical trials of several months to 2 years duration, symptomatic upper GI ulcers, gross bleeding or perforation occurred in approximately 1% of patients treated for 3 to 6 months, and in about 2% to 4% of patients treated for 1 year. Higher percentages have been reported by other independent studies. Studies to date have not identified any subset of patients not at risk of developing peptic ulceration and bleeding. Patients with prior history of serious GI events and other risk factors associated with peptic ulcer disease (e.g. alcoholism, smoking, corticosteroid therapy) are at increased risk. Elderly or debilitated patients seem to tolerate ulceration or bleeding less than other individuals and account for most spontaneous reports for fatal GI events. Measures such as the use of physical therapy and mild analgesics like paracetamol (when inflammation is not a major factor) should be instituted prior to initiation of therapy with NSAID. NSAIDs should only be used after proper appraisal of potential risks to patients. It should be used with the lowest effective dose for only as long as needed. This drug should not be co-administered with other NSAIDs. Prescribers should inform patients about the signs and/or symptoms of serious GI toxicity and what steps should take if

they occur. In considering the use of relatively large doses (within the recommended dosage range), sufficient benefit should offset the potential increased risk of GI toxicity.

USE IN PREGNANCY:

Ibuprofen is not recommended for use during pregnancy unless in the judgement of the physician the potential benefits to the mother outweigh the possible hazards to the foetus.

CONTRAINDICATIONS:

Known hypersensitivity to ibuprofen. Sensitive to aspirin or other non-steroidal anti-inflammatory drugs which includes those in whom attacks of asthma, angioedema, urticaria or rhinitis have been precipitated by aspirin or other NSAID. Active gastro-intestinal bleeding.

DRUG INTERACTIONS:

Non-steroidal anti-inflammatory drugs increase plasma concentration of lithium, methotrexate. It may reduce effects of diuretics, β -blockers. It may prolong bleeding time and increase plasma concentration of warfarin and may alter plasma concentration of phenytoin, sulfonyleureas (monitor).

RECOMMENDED DOSAGE:

Children:

1 ml/kg body weight daily given in divided doses.

6 months to 1 year as per above guide.

1 - 5 years old, 5 ml 3 times / day or 1 ml/kg body weight daily in divided doses.

5 - 12 years old, 10 ml 3 times a day or 1 ml/kg body weight daily in divided doses.

Up to 2 ml/kg body weight daily in divided doses may be given in cases of juvenile rheumatoid arthritis.

Adults:

20 ml, 4 to 6 hourly (maximum 120 ml a day).

SYMPTOMS AND TREATMENT FOR OVERDOSAGE:

Symptoms : Depression of CNS and respiratory system may occur.

Treatment : Evacuate stomach contents, institute general supportive measures and maintain urine output.

Alkalisiation of the urine may promote more rapid excretion.

EXPIRY PERIOD:

Do not use after the expiry date stated on the label.

STORAGE:

Keep below 30°C.

PACKING:

100 ml

DATE OF REVISION:

December 2021

For further information, please consult your physician or pharmacist.

Manufactured by:

PT. Actavis Indonesia
Jl. Raya Bogor Km. 28, Kelurahan Pekayon, Kecamatan Pasar Rebo Kota Jakarta Timur, Provinsi DKI Jakarta, Indonesia

Actavis

AAAN8431
40985-03