


ZEVIN CREAM
Aciclovir



COMPOSITION :
Each 100 gm contains : Aciclovir 5 g
DOSAGE FORMS AVAILABLE :
TOPICAL : CREAM 5%
PRODUCT DESCRIPTION :
A white cream with specific odor.
PHARMACODYNAMICS :
ZEVIN (Aciclovir) is a synthetic acyclic nucleoside analogue with in vitro inhibitory activity against herpes simplex types 1 and 2 (HSV-1 and HSV-2), varicella-zoster, Epstein-Barr and cytomegalovirus.
Mechanism of action: Aciclovir is converted to aciclovir monophosphate, a nucleotide, by herpes simplex virus (HSV)-coded thymidine kinase.
PHARMACOKINETICS :
Aciclovir is a synthetic nucleoside analogue antiviral drug. It is converted to aciclovir triphosphate, which is the pharmacological active form of the drug in infected cells.
INDICATION :
For the treatment of Herpes simplex virus infection of the skin including initial and recurrent genital herpes and herpes labialis.
DOSAGE AND ADMINISTRATION :
Apply ZEVIN CREAM five times daily at approximately four hourly intervals.
OVERDOSAGE, SYMPTOMS, TREATMENTS, SUPPORTIVE THERAPY, ANTIDOTE (S) :
Overdosage of aciclovir may cause renal dysfunction and eventually renal failure and anuria according to the precipitation of aciclovir crystals in the renal tubules.
CONTRAINDICATIONS :
ZEVIN is contraindicated in patients known to be hypersensitive to aciclovir or propylene glycol.
PRECAUTIONS AND WARNINGS :
1. Should avoid sexual contact while visible lesions are present because of the risk of infecting intimate partners.
2. Should be cautions not to exceed the prescribed dosage.
3. Do not apply to the mucous membrane eg. in the mouth, eyes or vagina as it may be irritant.
Use in pregnancy :
Limit data are available on the use of aciclovir in pregnancy.
Use in lactation :
Limited data indicate that aciclovir is distributed into milk following systemic administration.
ADVERSE EFFECTS :
Transient burning or stinging following application of aciclovir cream may occur.
DRUG INTERACTIONS :
Concomitant administration of probenecid and aciclovir has reportedly increased the mean plasma half life and area under the plasma concentration time curve (AUC) and decreased urinary excretion and renal clearance of aciclovir.
STORAGE CONDITIONS:
Store below 30°C. The shelf life period is 3 years.
SUPPLY :
Aluminium tube in printing box of 1x10 g
Date of revision : January 2019

Manufactured by
BIOLAB CO.,LTD.
625 Soi 7A Bangpoo Industrial Estate, Sukhumvit Road,
Moo 4, Prakasa, Muang, Samutprakarn 10280, Thailand

EER 019/B

230 mm.

100 mm.

Form containing Code, Draft Revision, Final Artwork Revision, Name, Remark, Design by, Review by, Approved by, Signature, Date, Specification, Size, Color, Material, Other, Note, COLOR GUIDE, and BIOLAB logo.