

DYNAMOL 650 TABLET

MAL19990615XZ

DESCRIPTION:

TABLET

Colour : Pink
Shape : Round, Flat, Scored
Marking : "DYNAMOL"

EACH TABLET CONTAINS:

Paracetamol.....650 mg

PHARMACODYNAMICS:

Paracetamol is considered to act mainly by inhibiting the biosynthesis of prostaglandins. It is often described as peripherally acting compound. Antipyretic activity is considered to involve the hypothalamus.

PHARMACOKINETICS:

Paracetamol is readily absorbed from the gastrointestinal tract with peak plasma concentrations occurring about 30 minutes to 2 hours after ingestion. It is metabolised in the liver and excreted in the urine mainly as the glucuronide and sulphate conjugates. Less than 5% is excreted as unchanged Paracetamol. The elimination half-life varies from about 1 hour to 4 hours. Plasma protein binding is negligible at usual therapeutic concentrations but increases with increasing concentrations. A minor hydroxylated metabolite which is usually produced in very small amounts by mixed-function oxidases in the liver and which is usually detoxified by conjugation with liver glutathione may accumulate following Paracetamol overdose and cause liver damage.

INDICATIONS:

Paracetamol has both analgesic and antipyretic action. It is used for the relief of pain such as headache, neuritis, myalgias and toothache, and fever.

RECOMMENDED DOSE:

Adults : 1 tablet.
Children 7 to 12 years : ½ tablet.
To be taken 3 to 4 times daily, as required.

ROUTE OF ADMINISTRATION:

FOR ORAL USE ONLY.

CONTRAINDICATIONS:

Hypersensitivity.

WARNINGS AND PRECAUTIONS:

This drug should be used with care in hepatic and renal impairment. Long term usage should be avoided. Do not exceed the recommended dose or use for more than 10 days unless prescribed.

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

This preparation contains PARACETAMOL.
Do not take any other PARACETAMOL containing medicines at the same time.

DRUG INTERACTION:

Paracetamol should not be taken concomitantly with any drug that acts on the liver.

PREGNANCY AND LACTATION:

Pregnancy: Not known to be harmful.
Lactation: Amount too small to be harmful.

SIDE EFFECTS:

Adverse effects are usually mild. Nausea, gastrointestinal upsets, rashes and allergic conditions have occurred. Overdose can cause hepatic damage, which is signified by jaundice, gastrointestinal bleeding and hypoglycaemia. Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome/Toxic Epidermal Necrolysis have been reported.

SYMPTOMS AND TREATMENT OF OVERDOSE:

Although the likely toxicity of a Paracetamol overdose can be assessed by examining the changing pattern of Paracetamol blood concentrations, prompt specific treatment is essential. Any patient therefore, who has taken 7.5 g or more of Paracetamol and can be treated within 10 hours of ingestion should be given a specific antidote, even if the blood concentrations are not known. There is a risk that the suphadryl donors used as antidotes may exacerbate any liver damage if given 10 hours after the overdose. Once specific therapy is established then blood concentrations of Paracetamol can be monitored. Generally treatment is required if the blood concentration is higher than a line (the '200' line) drawn on log/linear paper joining the points 200 mg per litre at 4 hours and 60 mg per litre at 10 hours. There have been suggestions that treatment may be necessary when lower concentrations of Paracetamol are present.

Cysteamine is sometimes used by injection as an antidote but acetylcystein or methionine is now preferred. Acetylcystein is given by intravenous infusion in an initial dose of 150 mg per kg body-weight over 15 minutes followed by 50 mg per kg over 4 hours and then 100 mg per kg over the next 16 hours. Alternatively, methionine 2.5 g may be given by mouth every 4 hours to a total of 4 doses. Haemoperfusion may be worthwhile if too much time has elapsed since the poisoning to allow use of acetylcystein or methionine. Basic measures that may be required include dextrose and blood infusions. Removal of stomach contents by aspiration and lavage forms an early part of treatment and the administration of Activated Charcoal should be considered.

PACKING/PACK SIZE(S):

Blister pack of 100x10's and 10x10's.

**JAUHI UBAT DARIPADA KANAK-KANAK
KEEP OUT OF REACH OF CHILDREN**

MANUFACTURER/PRODUCT REGISTRATION HOLDER:

DYNAPHARM (M) SDN. BHD. 198001011897 (65683-V)
2497, Mk. 1, Lorong Perusahaan Baru 5,
Kawasan Perusahaan Perai 3,
13600 Perai, Pulau Pinang, Malaysia.

PI3NT D030-04
Latest Revision: Jan 2022