



#### KETOZOL CREAM

Description: White opaque cream.

#### Content

Ketoconazole 2%w/w

Preservative: Chlorocresol 0.1%w/w

#### Indications

Pityriasis versicolor, cutaneous candidiasis, athlete's foot, tinea corporis, tinea cruris and seborrheic dermatitis.

#### Pharmacology

Ketoconazole is an imidazole antifungal agent which interferes with ergosterol synthesis and therefore alters the permeability of the cell membrane of sensitive fungi. It is reported to be fungistatic at concentration achieved clinically. Ketoconazole has a wide spectrum of antimicrobial activity in vitro including activity against *Candida sp.*, *Blastomycosis dermatitidis*, *Coccidioides immitis*, *Cryptococcus neoformans*, *Epidermophyton floccosum*, *Histoplasmosis capsulatum*, *Paracoccidioides brasiliensis*, *Malassezia furfur*, *Microsporum canis*, *Trichophyton mentagrophytes* and *T. rubrum*. Systemic absorption following topical application in healthy subjects is minimal.

#### Adverse effects

After topical application, irritation, dermatitis or a burning sensation has occurred.

#### Precautions

- Compliance with full course of therapy, fungal infection may require prolonged therapy
- Any topical steroid therapy should be reduced gradually when starting on topical ketoconazole cream to avoid skin irritation.
- This cream is not for ophthalmic use.
- Risk benefits have to be considered when administered on pregnant or lactating mothers.

#### Drug interactions:

Nil for topical application.

#### Contraindications

It is contraindicated in patients hypersensitive to ketoconazole.

#### Dosage and Administration

Apply once or twice daily and continue for at least a few days after the disappearance of symptoms. Depending on the severity of the infection, the usual duration of treatment for: pityriasis versicolor: 2-3 weeks, yeast infections: 2-3 weeks, tinea cruris: 2-4 weeks, tinea corporis: 3-4 weeks, tinea pedis: 4-6 weeks, seborrheic dermatitis: 2-4 weeks. Diagnosis should be reconsidered if no clinical improvement is noted after 4 weeks of treatment.

#### Treatment of overdosage

Prolonged use of topical application of ketoconazole may rarely lead to skin sensitization, resulting in hypersensitivity reactions with subsequent topical use of the medication. Discontinue use and wash with water the affected part.

Storage : Keep container tightly closed. Store in a cool, dry place below 30°C  
Protect from light. Keep out of reach of children.  
Presentation : 15gm/tube  
Shelf life : 3 years  
Manufactured & Distributed by : Prime Pharmaceutical Sdn. Bhd.  
1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah  
14000 Bukit Mertajam, Penang, Malaysia.

11/06/2019