

PORO[®] CHEWABLE TABLET 120MG

Ingredient:

Each chewable tablet contains:

Paracetamol (Acetaminophen) 120mg

Pharmacodynamics:

1. Paracetamol may act by inhibiting prostaglandin synthesis in the central nervous system and through a peripheral action by blocking pain impulse generation. The peripheral action may also be due to inhibition of the synthesis of prostaglandins which sensitize pain receptors to mechanical or chemical stimulation.
2. Paracetamol produces antipyresis by acting centrally on the hypothalamic heat-regulation center to produce peripheral vasodilation resulting in increase blood flow through the skin, sweating and heat loss. The central action probably involved inhibition of prostaglandin synthesis in the hypothalamus.

Pharmacokinetics:

Paracetamol is uniformly distributed throughout most body fluids, the apparent volume of distribution is 1 to 1.2 L/kg. Paracetamol can cross the placenta and is excreted in milk. At therapeutic plasma levels no protein binding occurs. Paracetamol is metabolised by the hepatic microsomal enzyme system. In adults at therapeutic doses, paracetamol is mainly conjugated with glucuronide (45-55%) or sulphate (20-30%). A minor proportion (less than 20%) is metabolised to catechol derivatives, and mercapturic acid compounds via oxidation. Paracetamol is metabolised differently by infants and children compared to adults, the sulphate conjugate being predominant. Paracetamol is excreted by active tubular secretion: 85-90% of the administered dose is eliminated in the urine within 24 hours of administration, Paracetamol has a half-life of 1 to 3.5 hours.

Indication(s):

Relieve symptoms of common cold, fever, headache, toothache and sore throat.

Dosage and Administration:

- 1-3 years : one tablet
4-6 years : one to two tablets,
7-12 years : two to four tablets.

Doses may be repeated every 4 to 6 hours when necessary.

Maximum of 4 doses in 24 hours.

Route of administration:

Oral administration.

Contraindication(s):

Contraindicated in patients who have shown hypersensitivity to Paracetamol.

Precaution(s) / Warning(s):

1. Use with caution in patient with renal or hepatic dysfunction.
2. If fever persists for more than 3 days or recurs, or if pain continues for more than 5 days, consult a physician immediately.
3. Allergy alert
Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash.
These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.
4. Unsuitable for phenylketonurics.

This preparation contains PARACETAMOL. Do not take any other paracetamol-containing medicines at the same time.

Drug Interaction:

Administration of paracetamol with Aspirin may increase blood concentrations of unhydrolysed Aspirin.

Side Effect(s) / Adverse Reaction(s):

Side effects of paracetamol are usually mild, though haematological reactions have been reported. Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome/Toxic Epidermal Necrolysis have been reported.

Symptoms and Treatment for Overdosage and Antidote(s):

Toxic symptoms include vomiting, abdominal pain, hypotension and sweating. Jaundice, hypoglycaemia and metabolic acidosis are major manifestation of liver failure, which may take at least 3 days to develop. In case of suspected overdose, getting emergency help at once even if no apparent symptoms. Symptoms of severe overdosage may be delayed, but treatment must be begun as soon as possible. Treatment started 24 hours or more after the overdosage may be ineffective in preventing liver damage or fatality. Acetylcysteine 20% is given by intravenous infusion as an antidote by a physician.

Shelf-Life:

3 years from the date of manufacture.

Storage Condition(s):

Store at temperature below 30°C. Protect from light and moisture.

Product Description and Packing(s):

Pinkish tablet with red dots scored on one side and impressed with "P" on the other side.

Blister packing of 10's x 2, 10's x 10, 10's x 50 and 10's x 100.

TABLET KUNYAH PORO®

Bahan Aktif:

Setiap tablet mengandung:
Paracetamol (Acetaminophen) 120mg

Indikasi:

Melegakan gejala selsema, demam, sakit kepala, sakit gigi dan sakit kerongkong.

Dos:

Kanak-kanak 1-3 tahun : satu tablet
Kanak-kanak 4-6 tahun : satu hingga dua tablet,
Kanak-kanak 7-12 tahun : dua hingga empat tablet.
Dos diulangi setiap 4 hingga 6 jam jika perlu.
Maksima 4 dos dalam 24 jam.

伯樂 甜嚼錠

成份:

每顆錠劑含:
Paracetamol (Acetaminophen) 120mg毫克

適應症:

解除感冒、發燒、頭痛、牙痛、喉嚨痛等疼痛。

用法、用量:

1 ~ 3 歲 : 1 粒。
4 ~ 6 歲 : 1 ~ 2 粒。
7 ~ 12 歲 : 2 ~ 4 粒。
以上劑量可每4至6 小時重复。在24小時內最多服用4次。



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679