



PARACETAMOL-250 SUSPENSION (STRAWBERRY)

COMPOSITION:

Each 5ml contains
Paracetamol 250mg

Methylparaben 0.1% w/v as preservative.

PRESENTATION:

Pinkish white suspension with strawberry flavour.

INDICATIONS:

For the relief of fever, symptoms of cold and influenza, teething pain and headache.

MECHANISMS OF ACTION:

Paracetamol is an antipyretic analgesic. The mechanism of action is probably similar to that of aspirin and dependent on the inhibition of prostaglandin synthesis. This inhibition appears, however to be on a selective basis.

PHARMACOLOGY:

Following oral administration paracetamol is rapidly absorbed. Paracetamol absorption takes place mainly in the small intestine and therefore the rate of absorption is depending on the rate of gastric emptying. It has been shown that drugs which delay gastric emptying also delay the absorption of paracetamol whereas metoclopramide (a drug which increases the rate of gastric emptying) accelerates absorption of the analgesic through the total amount absorbed doses not increase. The presence of food in the stomach has also been reported to reduce the rate of absorption of paracetamol. Alterations in gastric pH have no appreciable effect on paracetamol absorption. During absorption, the amount of paracetamol which is inactivated is negligible and it has been shown that paracetamol dose not affect gastric mucosal permeability and does not produce mucosal bleeding. Peak plasma concentrations are reached 1 hour after absorption. The plasma half-life is 1 to 3 hours. Paracetamol penetrates the brain and is present in breast milk of human. Paracetamol is metabolized by the microsomal enzyme system of the liver. This metabolism is mainly to the glucuronide and sulphate conjugates, accounting for approximately 49% and 26% of the ingested dose respectively. About 4% is excreted as free paracetamol. Other minor pathways include the production of catechol derivatives and cysteine conjugates (via glutathione). Paracetamol excretion is rapid and occurs via the urine.

DOSAGE AND ADMINISTRATION:

For oral administration only.
SHAKE WELL BEFORE USE.

Children:

1 - 6 years old : 2.5 - 5ml.

7 - 12 years old : 5 - 10ml.

To be taken 3 or 4 times a day.

Children 3 months to 1 year are advised to use Axcel Paracetamol-120 Syrup.

CONTRAINDICATION:

Contraindicated in patients with nephropathy.

PRECAUTIONS:

It should be given with care to patients with impaired kidney or liver functions. Chronic use should be avoided.

**THIS PREPARATION CONTAINS PARACETAMOL.
DO NOT TAKE ANY OTHER PARACETAMOL
CONTAINING MEDICINES AT THE SAME TIME.**

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Use in Pregnancy and Lactation:

Epidemiological studies in human pregnancy have shown no ill effects due to Paracetamol used in the recommended dosage, but patients should follow the advice of their doctor regarding its use. A large amount of data on pregnant women indicate neither malformative, nor foeto / neonatal toxicity. Epidemiological studies on neurodevelopment in children exposed to paracetamol in utero show inconclusive results. If clinically needed, paracetamol can be used during pregnancy however it should be used at the lowest effective dose for the shortest possible time and at the lowest possible frequency. Paracetamol is excreted in breast milk but not in a clinically significant amount. Available published data does not contraindicate breast-feeding.

SIDE EFFECTS:

Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome / Toxic Epiderma Necrolysis have been reported.

DRUG INTERACTION:

The prothrombinogenic effect of the oral anticoagulants may be increased by chronic administration of full doses of paracetamol.

OVERDOSAGE & TREATMENT :

Paracetamol overdose can result in severe liver damage and sometimes acute renal tubular necrosis. Immediate medical attention is essential in the event of overdose, even when there are no obvious symptoms. Administration of antidote such as acetylcysteine or methionine may be required. Activated charcoal may be used to reduce gastrointestinal absorption, if it can be given within 1 hour of the overdose, and if more than 150mg/kg of paracetamol has been ingested. However, acetylcysteine or methionine is to be given by mouth

the charcoal is best cleared from the stomach to prevent it reducing the absorption of the antidote.

STORAGE:

Keep container well closed. Store below 30°C. Protect from light.

**KEEP OUT OF REACH OF CHILDREN
JAUHI DARI KANAK-KANAK**

PACK QUANTITIES:

Available in PET bottle of 60ml & 100ml.

Further information can be obtained from doctor, pharmacist or the manufacturer.

MAL19991446XZ

Product Registration Holder and Manufacturer



Kotra Pharma (M) Sdn. Bhd.

No. 1, 2 & 3, Jalan TTC 12, (50052-V)

Cheng Industrial Estate,
75250 Melaka, Malaysia.

PMSPAXC05800 - 140322(03)
Date of revision: 24/10/23