

Uphamol 250



Product Description

Uphamol 250 Suspension 250mg/5ml (Fruity):
A pink, fruity flavoured suspension.

Uphamol 250 Suspension 250mg/5ml (Orange):
An orange, orange Flavoured suspension.

Active Ingredient:

Each 5ml (1 teaspoonful) contains:
Paracetamol (acetaminophen) 250mg

Preservatives:

Methylparaben 0.1% w/v

Pharmacodynamics

Analgesic

The mechanism of analgesic action has not been fully determined. Paracetamol may act predominantly by inhibiting prostaglandin synthesis in the central nervous system (CNS) and, to a lesser extent, through a peripheral action by blocking pain-impulse generation.

The peripheral action may also be due to inhibition of prostaglandin synthesis or to inhibition of the synthesis or actions of other substances that sensitize pain receptors to mechanical or chemical stimulation.

Antipyretic

Paracetamol probably produces antipyresis by acting centrally on the hypothalamic heat-regulating center to produce peripheral vasodilation resulting in increased blood flow through the skin, sweating, and heat loss. The central action probably involves inhibition of prostaglandin synthesis in the hypothalamus.

Pharmacokinetics

Paracetamol is rapidly and almost completely absorbed from the gastrointestinal tract following oral administration. Approximately 90-95% of a dose is metabolized in the liver, primarily by conjugation with glucuronic acid, sulfuric acid, and cysteine. An intermediate metabolite, which may accumulate in overdosage after the primary metabolic pathways become saturated, is hepatotoxic and possibly nephrotoxic. The half-life of paracetamol, 1 to 4 hours following oral administrations, does not change with renal failure but may be prolonged in acute overdosage, in some forms of hepatic disease, in the elderly, and in the neonate. However, it may be somewhat shortened in children. The time for paracetamol to reach its peak concentration is 0.5 to 2 hours and the time needed to reach the peak effect is 1 to 3 hours. Duration of action is 3 to 4 hours. It is excreted primarily as conjugates through renal and only 3% of a dose may be excreted unchanged.

Indication

Relieves fever, pain, discomforts of teething, headache, toothache, earache, common cold, influenza and post vaccination reactions.

Recommended Dosage

Dose to be taken every 4 to 6 hours up to a maximum of 4 doses in 24 hours.

1 to 5 years	½ to 1 teaspoonful (2.5 to 5 ml), 3 to 4 times daily
6 to 12 years	1 to 2 teaspoonful (5 to 10 ml), 3 to 4 times daily

Mode of Administration

Oral

Contraindications

Uphamol 250 should not be taken if children are allergic to acetaminophen (paracetamol)

Warnings and Precautions

If symptoms persist, please consult your doctor or pharmacist.

Do not keep outdated Uphamol 250 Suspension.

Allergy alert: Paracetamol may cause severe skin reactions.

Symptoms may include skin reddening, blisters or rash. These

could be signs of a serious condition.

If these reactions occur, stop use and seek medical assistance right away.

This preparation contains PARACETAMOL.
Do not take any paracetamol
containing medicines at the same time.

Adverse Effects / Undesirable Effects

When taken as directed, unwanted effects from Uphamol 250 are rare and usually mild. Skin rashes and other hypersensitivity reactions have been reported occasionally.

Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome/Toxic Epidermal Necrolysis have been reported.

Overdose and Treatment

Symptoms :

Vomiting, mild nausea, gastrointestinal upset, hypotension, anorexia and diarrhoea.

Treatment (for medical practitioner's information) :

- Empty the stomach via induction of emesis or gastric lavage.
- Acetylcysteine administration is recommended to be instituted immediately after ingestion of an overdose has been reported without waiting for the result of urine and plasma level tests. Acetylcysteine is most effective if treatment is started within 10 to 12 hours after ingestion of overdose, however, it may be some benefit if treatment is started within 24 hours. For oral administration, the recommended adult dose of acetylcysteine is 140mg/kg initially, then 70mg/kg every 4 hours for 17 doses. Each dose should be diluted to a 5% solution of drinking fluid prior to administration because of its unpleasant odour and its sclerosing properties. Any dose vomited within 1 hour of administration must be repeated. If necessary, the antidote may be given via duodenal intubation.

- Plasma acetaminophen concentration should be determined at least 4 hours following the ingestion of overdose in order to ascertain the status of patient. Renal and cardiac function should be monitored and appropriate therapy instituted if necessary.

Storage Conditions

Store in a dry place below 30°C.

Keep container tightly closed.

Protect from light.

Keep out of reach of children.

Jauh dari kanak-kanak

Pack Size

60ml, 100ml, 72 x 60ml and 72 x 100ml

Shelf Life

Please refer to box or label.



Product Registration Holder:

DUOPHARMA MANUFACTURING (BANGI) SDN. BHD.

Lot No. 2,4,6,8 & 10, Jalan P/7, Section 13, Bangi Industrial Estate,
43650 Bandar Baru Bangi, Selangor, Malaysia.

Manufacturer:

DUOPHARMA MANUFACTURING (BANGI) SDN. BHD.

Lot No. 2 & 4, Jalan P/7, Section 13, Bangi Industrial Estate,
43650 Bandar Baru Bangi, Selangor, Malaysia

Date of Revision

December 2025

1500008810 N.1

Size: 23.2cm (H) x 15cm (W)



Black 100%