

KEZOL CREAM

DESCRIPTION:

A white, smooth homogeneous mass

COMPOSITION:

Each gm of the cream contains Ketoconazole (Micronised) 20mg

ACTIONS & PHARMACOLOGY:

Ketoconazole has a potent antimycotic against dermatophytes. It is fungistatic; may be fungicidal, depending on concentration; inhibits biosynthesis of ergosterol or other sterols, damaging the fungal cell membrane and altering its permeability; as a result, loss of essential intracellular elements may occur, also inhibits biosynthesis of triglycerides and phospholipids by fungi; in addition, inhibits oxidative and peroxidative enzyme activity, resulting in intracellular buildup of toxic concentrations of hydrogen peroxide, which may contribute to deterioration of subcellular organelles and cellular necrosis. In the treatment of *Candida albicans*, inhibits transformation of blastospores into invasive mycelial form. Systemic absorption following topical application to normal skin has not been reported.

INDICATIONS:

Dermatophyte infections of the skin; Tinea corporis, Tinea cruris, Tinea manus and Tinea pedis due to *Trichophyton rubrum*, *T. mentagrophytes*, *Microsporum canis* and *Epidermophyton floccosum*; as well as in the treatment of cutaneous candidosis and tinea (pityriasis) versicolor. Seborrhoeic dermatitis, a skin condition related with the presence of *Pityrosporum ovale*.

CONTRAINDICATIONS:

Contraindicated in patients who have shown hypersensitivity to any of the ingredients and Ketoconazole.

SIDE EFFECTS:

Few instances of irritation and burning sensation have been observed during treatment with the cream. Irritation may occur when cream is used immediately after prolonged treatment with topical corticosteroids.

DRUG INTERACTION:

None Known

WARNING AND PRECAUTIONS:

Prolonged use of topical ketoconazole may rarely lead to skin sensitization, resulting in hypersensitivity reaction with subsequent topical or systemic use of the medication. To reduce the possibility of recurrence, comply with full course of therapy. Use hygienic measures to cure infection and prevent reinfection. Avoid contact with the eyes.

Recovery periods of two weeks should be allowed, if potent corticosteroid has been used to treat seborrhoeic dermatitis. This product contains Sodium sulphite that may cause serious allergic type reactions in certain susceptible patients.

Children: Safety and efficacy have not been established

PREGNANCY AND LACTATION:

Ketoconazole crosses the placenta. Adequate and well-controlled studies in humans have not been done. Topical ketoconazole is unlikely to be distributed into breast milk in significant amounts or to cause adverse effects in the nursing infant.

DOSAGE & ADMINISTRATION:

Apply to the affected area once daily in patients with Tinea corporis, Tinea cruris, Tinea pedis, Pityriasis versicolor and Cutaneous candidiasis.

Twice daily in patients with seborrhoeic dermatitis.

Treatment should be continued for a sufficient period, at least a few days disappearance of all symptoms. Diagnosis should be reconsidered if no clinical improvement is noted after 4 weeks.

ROUTE OF ADMINISTRATION

Topical administration

EFFECTS ON ABILITY TO DRIVE AND USE MACHINE

Not Applicable

OVERDOSAGE & TREATMENT:

Acute overdosage with the topical application of the drug is unlikely. If irritation or sensitivity reactions occur, use of the medication should be discontinued.

PRESENTATIONS:

Aluminium tubes of 5g, 10g, 15g, and 50g.

STORAGE CONDITIONS:

Store below 30°C. Keep the containers tightly closed.

For external use only. Keep medicine out of reach of children. Protect from light and freezing.

MANUFACTURER BY:

NORIPHARMA SDN. BHD. 200701034604 (792633-A)

Lot 5030, Jalan Teratai,

5 ½ Mile off Jalan Meru

41050 Klang,

Selangor Darul Ehsan., Malaysia

DATE OF REVISION:

May 2021