



Ubat Serbuk Cap Ikan Emas Aspirin 600mg/Paket

Gold Fish Trade Mark Medicinal Powder

KOMPOSISI : Setiap paket mengandungi Aspirin 600mg.

DESKRIPSI : Serbuk yang berwarna putih, tidak berbau atau hampir tidak berbau dengan sedikit rasa asid.

INDIKASI : Untuk melegakan demam, tanda selsema, sakit kepala, sakit gigi dan sakit otot.

SUKATAN DAN PENGGUNAAN : 1) Sakit kepala, demam, sakit gigi, sakit otot, sakit haid, migraine : 1 pekset setiap 4 jam jika perlu. Maksima 6 kali sehari. 2) : Untuk keadaan keradangan dan sakit arthritis dan rheumatoid arthritis : 2 pekset setiap 6 jam jika perlu. maksima 4 kali sehari. Mesti diambil selepas makan. Campur dengan segelas air sebelum makan.

PERHATIAN/AMARAN : Aspirin mesti dibagi dengan perhatian kepada pesakit yang mengidap kelukaan gastrik mukosa atau keadaan terjejas renal atau hepatic. Tidak boleh diberi kepada kanak-kanak berumur kurang daripada 16 tahun. Kegunaan harian mungkin akan mengakibatkan antikoagulan lisan dan agen sulphonylurea hipoglisemik.

KESAN SAMPINGAN : Kebiasaan adalah gangguan radang, keususan perut seperti rasa loya, dispepsia dan muntah. Kemungkinan berlaku kerangsangan pada bendalir perut dan ulser. Aspirin akan meningkatkan masa pendarahan dan menurunkan tahap kelekatan platelet.

KONTRAINDIKASI : Tidak boleh diambil oleh pesakit yang mengidap penyakit haemophilia, kerosakan hati, kerosakan buah pinggang, ulser gastrik dan alergi kepada aspirin.

SIMPON UNTUK KELEBIHAN DOS :
Rujuk penerangan Bahasa Inggeris.

RAWATAN UNTUK KELEBIHAN DOS : Rujuk penerangan Bahasa Inggeris.

INTERAKSI : Rujuk penerangan Bahasa Inggeris.

FARMAKODINAMIK : Rujuk penerangan Bahasa Inggeris.

FARMAKOKINETIK : Rujuk penerangan Bahasa Inggeris.

PENYIMPANAN : Simpan di bawah 30°C. Jauhi dari kanak-kanak.

TEMPOH HAYAT : Jangan guna selepas tarikh luput.

KANDUNGAN : Satu kotak mengandungi 12 paket. Berat bersih setiap paket : 0.95gm.

****Maklumat yang terkandung dalam surat ini adalah terhad. Untuk maklumat yang terperinci, sila hubungi agen tempatan, doktor atau ahli farmasi.**

Dibuat oleh/Manufactured by:
SYARIKAT WEN KEN DRUG SDN BHD
24, Jalan Lambak, Taman Johor, 81200 Johor Bahru, Johor, Malaysia.

Manufactured for Product Registration Holder:
ALPINE MEDICAL INDUSTRIES SDN. BHD.
34, Jalan Perindustrian Silibin 1, Kawasan Perindustrian Ringan Silibin, 30100 Ipoh, Perak.

Revision No.: 03
Revision date : 20 August 2018

COMPOSITION : Each sachet contains Aspirin 600mg.

DESCRIPTION: White powder, odourless or almost odourless with a slight acid taste.

INDICATIONS : For the relief of fever and discomfort associated with colds, headache, toothache and muscular pains.

DOSAGE AND ADMINISTRATION : 1) Headache, fever, toothache, muscle pain, menstrual pain, migraine : 1 sachet every 4 hours if required. Maximum 6 times a day. 2) for inflammatory condition and pain due to arthritis and rheumatoid arthritis : 2 sachets every 6 hours if required. Maximum 4 times a day. The dose should be taken after meals. Dissolve the medicinal powder in a glass of water before taking.

PRECAUTIONS/WARNINGS : Aspirin should be given with care to patient known to have lesion of the gastric mucosa or patients with impaired renal or hepatic function. Not to be given to children under 16 years of age. Daily use may potentiate oral anticoagulants and sulphonylurea hypoglycaemic agents.

SIDE EFFECTS : The most common are gastro-intestinal disturbances such as nausea, dyspepsia and vomiting. Irritation of the gastric mucosa and ulceration may occur. Aspirin increase the bleeding time and decrease platelet adhesiveness.

CONTRAINDICATIONS : It should not be administered to patient with haemophilia. It is contraindicated in severe impaired renal and hepatic function, ulceration and known allergy to aspirin.

SYMPTOMS OF OVERDOSE : Dizziness, tinnitus, deafness, sweating, nausea and vomiting, headache & mental confusion. In severe poisoning, hyperventilation, fever, restlessness, ketosis and respiratory alkalosis and metabolic acidosis result. Depression of CNS may lead to coma, cardiovascular collapse and respiratory failure.

TREATMENT OF OVERDOSE : In acute salicylate overdose, the stomach should be emptied by including emesis or by aspiration and lavage. Patients with mild intoxication should be encouraged to drink plenty of fluids. In patients with more severe intoxication (plasma salicylate concentration above 500µg per ml in adults or 300µg per ml in children), forced alkaline diuresis maybe required and should be continued until the plasma salicylate concentration is less than 350µg per ml in adults.

DRUG INTERACTION : Alcohol, coumarin, anticoagulants, sulphonylurea hypoglycaemic agents, methotrexate, uricosuric agents, barbiturates and other sedatives.

PHARMACODYNAMICS : Aspirin has analgesic and antipyretic as well as anti-inflammatory effects. It alleviates pain by inhibiting prostaglandin synthesis in the central nervous system and through a peripheral action by inhibiting the syththesis of prostaglandins that occurs in inflamed tissues to prevent the sensitization of the pain receptors to mechanical or chemical stimulation. Aspirin produces antipyresis by acting centrally on the hypothalamic heat-regulating center by inhibiting prostaglandin synthesis in the hypothalamus to produce peripheral vasodilation, resulting in increased cutaneous blood flow, sweating and heat loss. Its anti-inflammatory activity is by inhibiting the synthesis of prostaglandin in the inflamed tissue.

PHARMACKINETICS : Absorption is rapid and complete following oral administration mostly from the upper small intestine. Food decrease the rate but not the extent of absorption. The plasma concentration reaches a peak in 1 to 2 hours, and the plasma half-life is about 2 to 3 hours Aspirin is primarily metabolized in the liver. It is excreted in the urine as salicylic acid (10%) and as glucuronide conjugates (15%) and as salicyluric (75%) and gesticic acids (<1%). The rate of excretion of aspirin varies with the pH of the urine increasing as the pH rises.


STORAGE : Store below 30°C
Keep out of reach of children.

SHELF LIFE : Do not use beyond expiry date.

PACKING : Each box contains 12 sachet.
Net weight of each sachet : 0.95gm.

****The information contained in this leaflet is limited. For further information, please consult your doctor, pharmacist or the local agent.**

210mm



金鱼标外感散阿司匹林600毫克/包

成份 : 每包含有600毫克 Aspirin。

外观 : 无味，带点酸性的白色药粉。

适应症 : 能有效的解除发烧，伤风，头痛，牙痛及肌肉疼痛。

服量 : 成人每次服一包，将整包溶于水中，若需要的话，可每隔四小时再服用，每日最多服用6次，若症状仍持续，应请教医生。

防范 : 对于患上消化不良，消化性溃疡或敏感性症候（特别是哮喘）的患者，则应在医药护理人员的监督之下才可服用。不可服食超过以上服量。

副作用 : 可能产生消化不良，作呕或呕吐现象，严重的可能引起溃疡及粪便带血，要避免上述情形，应在饭后服用。

禁忌 : 此药粉不可给与血友病患者服用以及一些肝功能有损，胃溃疡或对阿司匹灵敏感者。

服过量症状及医治 : 请参考英文说明。

药物相互作用 : 请参考英文说明。


储藏 : 应置放於普通室温及儿童无法取到处。

注意 : 十六岁以下儿童不宜服用，除非得到医生的指示。

有效期 : 不要使用超出有效时期。

包装 : 每盒含有12包粉囊。
每包粉囊净重0.95克。

****此说明书仅属略述，欲知更详细资料，可询问您的医生，药剂师或邻近代理商。**



Ubat Serbuk Cap Ikan Emas Aspirin 600mg/Paket

Gold Fish Trade Mark Medicinal Powder

KOMPOSISI : Setiap paket mengandungi Aspirin 600mg.

DESKRIPSI : Serbuk yang berwarna putih, tidak berbau atau hampir tidak berbau dengan sedikit rasa asid.

INDIKASI : Untuk melegakan demam, tanda selsema, sakit kepala, sakit gigi dan sakit otot.

SUKATAN DAN PENGGUNAAN : 1) Sakit kepala, demam, sakit gigi, sakit otot, sakit haid, migraine : 1 pekset setiap 4 jam jika perlu. Maksima 6 kali sehari. 2) : Untuk keadaan keradangan dan sakit arthritis dan rheumatoid arthritis : 2 pekset setiap 6 jam jika perlu. maksima 4 kali sehari. Mesti diambil selepas makan. Campur dengan segelas air sebelum makan.

PERHATIAN/AMARAN : Aspirin mesti dibagi dengan perhatian kepada pesakit yang mengidap kelukaan gastrik mukosa atau keadaan terjejas renal atau hepatic. Tidak boleh diberi kepada kanak-kanak berumur kurang daripada 16 tahun. Kegunaan harian mungkin akan mengakibatkan anticoagulan lisan dan agen sulphonylurea hipoglisemik.

KESAN SAMPINGAN : Kebiasaan adalah gangguan radang, keususan perut seperti rasa loya, dispepsia dan muntah. Kemungkinan berlaku kerangsangan pada bendalir perut dan ulser. Aspirin akan meningkatkan masa pendarahan dan menurunkan tahap kelekatan platelet.

KONTRAINDIKASI : Tidak boleh diambil oleh pesakit yang mengidap penyakit haemophilia, kerosakan hati, kerosakan buah pinggang, ulser gastrik dan alergi kepada aspirin.

SIMPON UNTUK KELEBIHAN DOS : Rujuk penerangan Bahasa Inggeris.

RAWATAN UNTUK KELEBIHAN DOS : Rujuk penerangan Bahasa Inggeris.

INTERAKSI : Rujuk penerangan Bahasa Inggeris.

FARMAKODINAMIK : Rujuk penerangan Bahasa Inggeris.

FARMAKOKINETIK : Rujuk penerangan Bahasa Inggeris.

PENYIMPANAN : Simpan di bawah 30 °C. Jauhi dari kanak-kanak.

TEMPOH HAYAT : Jangan guna selepas tarikh luput.

KANDUNGAN : Satu kotak mengandungi 12 paket. Berat bersih setiap paket : 0.95gm.

****Maklumat yang terkandung dalam surat ini adalah terhad. Untuk maklumat yang terperinci sila hubungi agen tempatan, doktor atau ahli farmasi.**

Dibuat oleh/Manufactured by:
SYARIKAT WEN KEN DRUG SDN BHD
24, Jalan Lambak, Taman Johor, 81200 Johor Bahru, Johor, Malaysia.

Manufactured for Product Registration Holder:
ALPINE MEDICAL INDUSTRIES SDN. BHD.
34, Jalan Perindustrian Silibin 1, Kawasan Perindustrian Ringan Silibin, 30100 Ipoh, Perak.

Revision No.: 03
Revision date : 20 August 2018

COMPOSITION : Each sachet contains Aspirin 600mg.

DESCRIPTION: White powder, odourless or almost odourless with a slight acid taste.

INDICATIONS : For the relief of fever and discomfort associated with colds, headache, toothache and muscular pains.

DOSAGE AND ADMINISTRATION : 1) Headache, fever, toothache, muscle pain, menstrual pain, migraine : 1 sachet every 4 hours if required. Maximum 6 times a day. 2) for inflammatory condition and pain due to arthritis and rheumatoid arthritis : 2 sachets every 6 hours if required. Maximum 4 times a day. The dose should be taken after meals. Dissolve the medicinal powder in a glass of water before taking.

PRECAUTIONS/WARNINGS : Aspirin should be given with care to patient known to have lesion of the gastric mucosa or patients with impaired renal or hepatic function. Not to be given to children under 16 years of age. Daily use may potentiate oral anticoagulants and sulphonylurea hypoglycaemic agents.

SIDE EFFECTS : The most common are gastro-intestinal disturbances such as nausea, dyspepsia and vomiting. Irritation of the gastric mucosa and ulceration may occur. Aspirin increase the bleeding time and decrease platelet adhesiveness.

CONTRAINDICATIONS : It should not be administered to patient with haemophilia. It is contraindicated in severe impaired renal and hepatic function, ulceration and known allergy to aspirin.

SYMPTOMS OF OVERDOSE : Dizziness, tinnitus, deafness, sweating, nausea and vomiting, headache & mental confusion. In severe poisoning, hyperventilation, fever, restlessness, ketosis and respiratory alkalosis and ~~metabolic acidosis result. Depression of CNS may lead to coma,~~ cardiovascular collapse and respiratory failure.

TREATMENT OF OVERDOSE : In acute salicylate overdose, the stomach should be emptied by including emesis or by aspiration and lavage. Patients with mild intoxication should be encouraged to drink plenty of fluids. In patients with more severe intoxication (plasma salicylate concentration above 500µg per ml in adults or 300µg per ml in children), forced alkaline diuresis maybe required and should be continued until the plasma salicylate concentration is less than 350µg per ml in adults.

DRUG INTERACTION : Alcohol, coumarin, anticoagulants, sulphonylurea hypoglycaemic agents, methotrexate, uricosuric agents, barbiturates and other sedatives.

PHARMACODYNAMICS : Aspirin has analgesic and antipyretic as well as anti-inflammatory effects. It alleviates pain by inhibiting prostaglandin synthesis in the central nervous system and through a peripheral action by inhibiting the sythesis of prostaglandins that occurs in inflamed tissues to prevent the sensitization of the pain receptors to mechanical or chemical stimulation. Aspirin produces antipyresis by acting centrally on the hypothalamic heat-regulating center by inhibiting prostaglandin synthesis in the hypothalamus to produce peripheral vasodilation, resulting in increased cutaneous blood flow, sweating and heat loss. Its anti-inflammatory activity is by inhibiting the synthesis of prostaglandin in the inflamed tissue.

PHARMACKINETICS : Absorption is rapid and complete following oral administration mostly from the upper small intestine. Food decrease the rate but not the extent of absorption. The plasma concentration reaches a peak in 1 to 2 hours, and the plasma half-life is about 2 to 3 hours Aspirin is primarily metabolized in the liver. It is excreted in the urine as salicylic acid (10%) and as glucuronide conjugates (15%) and as salicyluric (75%) and gestisic acids (<1%). The rate of excretion of aspirin varies with the pH of the urine increasing as the pH rises.

STORAGE : Store below 30°C
Keep out of reach of children.

SHELF LIFE : Do not use beyond expiry date.

PACKING : Each box contains 12 sachet. Net weight of each sachet : 0.95gm.

****The information contained in this leaflet is limited. For further information, please consult your doctor, pharmacist or the local agent.**

210mm



金鱼标外感散阿司匹林600毫克/包

成份 : 每包含有600毫克 Aspirin。

外观 : 无味，带点酸性的白色药粉。

适应症 : 能有效的解除发烧，伤风，头痛，牙痛及肌肉疼痛。

服量 : 成人每次服一包，将整包溶于水中，若需要的话，可每隔四小时再服用，每日最多服用6次，若症状仍持续，应请教医生。

防范 : 对於患上消化不良，消化性溃疡或敏感性症候（特别是哮喘）的患者，则应在医药护理人员的监督之下才可服用。不可服食超过以上服量。

副作用 : ~~可能产生消化不良，作呕或呕吐现象，严重的可能引起溃疡及粪便带血，要避免上述情形，应在饭后服用。~~

禁忌 : 此药粉不可给与血友病患者服用以及一些肝功能有损，胃溃疡或对阿司匹灵敏感者。

服过量症状及医治 : 请参考英文说明。

药物相互作用 : 请参考英文说明。

储藏 : 应置放於普通室温及儿童无法取到处。

注意 : ~~十六岁以下儿童不宜服用，除非得到医生的指示。~~

有效期 : 不要使用超出有效时期。

包装 : 每盒含有12包粉囊。每包粉囊净重0.95克。

****此说明书仅属略述，欲知更详细资料，可询问您的医生，药剂师或邻近代理商。**

148mm

Remarks:
1. Updated statement as per latest DRGD
2. Updated statement as per latest DRGD
3.