

PORO[®] SUSPENSION 250MG/5ML (STRAWBERRY FLAVOUR)

Each 5ml contains:

Ingredient(s):

Paracetamol 250mg

Preservatives:

Methyl Paraben 0.18%w/v

Propyl Paraben 0.02%w/v

Pharmacology (Summary of Pharmacodynamics and Pharmacokinetics):

The analgesic action of Paracetamol involves peripheral and central influences. Antipyretic is mediated through hypothalamic heat regulating centers. Paracetamol inhibits prostaglandin synthetase. Therapeutic doses of Paracetamol have negligible effects on the cardiovascular respiratory systems; however, toxic doses may cause circulatory failure and rapid, shallow breathing.

Indication(s):

For the relief of fever and mild to moderate pain.

Dosage and Administration:

1 ~ 5 years : 2.5 ml ~ 5 ml ($\frac{1}{2}$ ~ 1 teaspoon)

6 ~ 12 years : 5ml ~ 10ml (1 ~ 2 teaspoon)

Doses may be repeated every 4 ~ 6 hours when necessary.

(max of 4 doses in 24 hours).

To be taken orally.

Shake well before use

Contraindications:

Hypersensitivity to Paracetamol.

Precaution(s) / Warning(s):

If fever persists for more than 3 days or recurs, or if pain continue for more than 5 days, consult a physician immediately. Use with caution in patients with anemia or with cardiac, pulmonary, renal or hepatic disease.

Allergy alert:

Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash.

These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

This preparation contains PARACETAMOL. Do not take any other PARACETAMOL containing medicines at the same time.

Drug Interaction:

1. Concurrent chronic, high-dose administration of Paracetamol with anticoagulants may increase the anticoagulant effect.
2. Administration of Paracetamol with Aspirin increases blood concentration of unhydrolysed aspirin.

Pregnancy and Lactation

Pharmaceutical studies in 12 nursing mothers given a single dose of Paracetamol suggested that maternal ingestion of Paracetamol in usual therapeutic doses does not appear to present a risk to nursing infant.

Paracetamol metabolites were found in the neonate following maternal overdose at 36 weeks. The baby was delivered by Caesarean section and did not suffer liver damage; the mother was treated with Acetylcysteine.

Side Effect(s) / Adverse Reaction(s):

Side effects of Paracetamol are usually mild, though hematological reactions have been reported. Skin rashes and other allergic reactions occur occasionally.

Cutaneous hypersensitivity reactions including skin rashes, angioedema, Steven-Johnson Syndrome/Toxic Epidermal Necrolysis have been reported.

Symptoms and Treatment For Overdosage, and Antidote(s):

Initial symptoms of overdosage include nausea, vomiting, hypotension and sweating. Liver damage is unlikely to occur after ingestion of less than 15g Paracetamol (300mL) as a single dose. Jaundice, hypoglycaemia and metabolic acidosis are the major manifestations of liver failure which may take at least three days to develop. In cases of suspected overdose, Acetylcysteine is given by intravenous infusion as an antidote by a physician.

Storage Condition(s):

Keep in an airtight container. Store at temperature below 30°C. Protect from light and moisture. Keep medicine out reach of children.

Shelf-Life:

3 years from the date of manufacture.

Product description and Packing(s):

Pink color, strawberry flavored suspension.

Plastic bottles of 60ml, 100ml.

SUSPENSI PORO® 250MG/5ML (PERISA STRAWBERI)

Setiap 5ml mengandungi:

Bahan Aktif:

Paracetamol 250mg

Bahan pengawet:

Methyl Paraben 0.18%w/v

Propyl Paraben 0.02%w/v

Indikasi:

Untuk melegakan kesakitan yang ringan dan sederhana serta mengurangkan demam.

Dos:

Kanak-kanak :

1 ~ 5 tahun : 2.5ml ~ 5ml ($\frac{1}{2}$ ~ 1 camca teh)

6 ~ 12 tahun : 5ml ~ 10ml (1~2 camca teh)

Dos boleh diulang setiap 4 - 6 jam jika perlu sahaja.

(maksimum 4 dos dalam 24 jam).

Goncang sebelum minum

伯樂懸液 250MG/5ML (草莓味)

每5ml含:

成分:

Paracetamol 250mg

防腐劑:

Methyl Paraben 0.18%w/v

Propyl Paraben 0.02%w/v

適應症:

1. 伯樂懸液是一種溫和的止痛劑，其主要成分PARACETAMOL能夠提升疼痛的閾值，適用於小兒感冒引起的頭痛。
2. 本劑亦作用在下視丘之體溫控制中樞，促進皮膚層的血流量，造成出汗及解熱。
3. 伯樂懸液適口，小兒易服用。

劑量:

年齡(歲)	1~5歲	6~12歲
劑量 (茶匙)	2.5~5ml ($\frac{1}{2}$ ~ 1)	5~10ml (1~2)

每4~6小時服用一次，一天不能超過4次。



Manufacturer and Product Registration Holder:

Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)

Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.

Ordering Line: 1 800 88 3027

Product Info: 1 800 88 3679