



GENTASONE CREAM

Description: White colour cream.

Use of topical gentamicin in closed hospital settings is actively discouraged.

Content:

Gentamicin sulphate eq. to gentamicin 0.10%w/w
Betamethasone dipropionate eq. to betamethasone 0.05%w/w

Indication:

Dermatoses complicated by or when bacterial infection is suspected.

Pharmacology:

Gentamicin sulphate is a broad spectrum aminoglycoside antibiotic which is effective against a wide range of Gram +ve and Gram -ve bacteria, including *Staphylococcus aureus* (coagulase +ve, coagulase -ve and some penicillinase-producing strains), *Streptococcus* group A (α and β hemolytic), species of *Pseudomonas*, *Proteus vulgaris*, *E. coli*, *Aerobacter aerogenes*, *Klebsiella pneumoniae* and many coliform bacilli. It is bactericidal and acts by binding to a specific receptor protein on the 30S subunit inhibiting protein synthesis. DNA may be misread, thus producing non-functional proteins, polyribosomes are split apart and unable to synthesize proteins. Betamethasone dipropionate is a glucocorticoid applied topically in the management of various disorders in which corticosteroids are indicated. It has marked antipruritic, vasoconstrictor and anti-inflammatory activity.

Systemic absorption of gentamicin has been reported after topical use on denuded skin and burns. Betamethasone dipropionate is well absorbed from the sites of local application, particularly under an occlusive dressing or when the skin is broken. It may be absorbed to give systemic effects. Metabolic reactions of gentamicin is by conjugation. Binding of gentamicin to plasma proteins is usually low. Betamethasone dipropionate is more slowly metabolized in the skin and this tend to be systemically absorbed in a greater extent.

Adverse effects:

Topical application of gentamicin may give rise to allergy e.g. redness, swelling, itching or other signs of irritation not present before therapy. When applied topically, particularly to large areas when the skin is broken or under occlusive dressings, betamethasone dipropionate may be absorbed in sufficient amounts to cause systemic effects. With prolonged use, maceration of the skin, skin atrophy, striae and secondary infection may be present.

Precautions:

Use of topical antibiotics occasionally allows overgrowth of the nonsusceptible organisms, including fungi. If this occurs or if irritation, sensitization or superinfection developed, treatment with gentamicin should be discontinued and appropriate therapy instituted. Steroids applied topically over large areas, particularly when the skin is broken, with or without occlusive dressings may result in systemic effects. Therefore, suitable precautions should be taken. Special precautions has to be observed when use on infants and children less than 4 years as it may cause growth retardation on prolonged use. Therefore, avoid prolonged use in children and on the face.

Pregnancy & Lactation:

Risk benefits should be considered when use in pregnant and lactating mothers, although safety of use has not been established.

Drug Interactions:

Nil for topical application.

Contraindications:

It is contraindicated in patients hypersensitive to gentamicin, betamethasone dipropionate or any related components, in vaccinia, varicella and tuberculosis of the skin.

Dosage and administration:

Apply thinly to affected skin two times daily or less frequent.

Treatment of overdosage:

Excessive or prolonged use of topical corticosteroids can suppress pituitary-adrenal function, resulting in secondary adrenal insufficiency, and produce manifestations of hypercorticism, including Cushing's disease. Excessive or prolonged use of gentamicin may cause overgrowth of lesions by fungi or nonsusceptible bacteria. In cases of chronic toxicity, slow withdrawal of steroids is advised. Appropriate antifungal or antibacterial therapy is indicated if overgrowth occurs. Symptomatic treatment may be applied.

Storage conditions : Keep container tightly closed. Store in a cool, dry place below 30C. Protect from light. Keep away from children.

Presentation

: 15gm tube

Shelflife

: 3 years

Manufactured &

Distributed by

PRIME PHARMACEUTICAL SDN. BHD.

1505, Lorong Perusahaan Utama 1,

Taman Perindustrian Bukit Tengah,

14000 Bukit Mertajam, Penang, Malaysia.