

**AMMI VOTARA EMULSION GEL**

|                                      |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
|--------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Description</b>                   | : Smooth, white water soluble gel with characteristic odour.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| <b>Composition</b>                   | : Diclofenac Diethylamine Salt 1.16% w/w<br>equivalent to Diclofenac Sodium 1.0% w/w                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| <b>Pharmacodynamics</b>              | : Diclofenac has been shown in experiment to inhibit prostaglandin biosynthesis, it is the important mechanism of action. Ammi Votara Emulsion Gel is prepared for external application as an analgesic and anti-inflammatory gel, easily be rubbed into the skin with a soothing and cooling effect, relieves pain, reduces oedema and shortens the time to return normal function.                                                                                                                                                                                                                                                                                                                                                                                      |
| <b>Pharmacokinetics</b>              |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Absorption                           | : The amount of diclofenac absorbed through the skin is proportional to the contact time and skin area covered and depends on the total topical dose and on skin hydration.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| Distribution                         | : Diclofenac can be detected in plasma, synovial tissue and fluid after topical application.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Metabolism & Elimination             | : It is excreted mainly in the urine. No accumulation of diclofenac and its metabolites should occur in patients with renal insufficiency. In patients with chronic hepatitis or nondecompensated liver cirrhosis, the kinetics and metabolism of diclofenac are the same as in patients without liver disease.                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| <b>Indications</b>                   | : Symptomatic relief and treatment of soft-tissue rheumatism, e.g. tendovaginitis, shoulder-hand syndrome, bursitis and periarthopathy; rheumatic diseases, e.g. osteoarthritis of the spine and peripheral joints; post-traumatic inflammation of the tendons, ligaments, muscles and joints. e.g. due to sprains, strains or bruises.                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| <b>Route of Administration</b>       | : Topical                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| <b>Recommended Dosage</b>            | : After assessing the risk/benefit ratio in each individual patient, the lowest effective dose for the shortest possible duration should be used.<br>Apply for 2-4 g, 3-4 times daily to affected sites and rub-in gently. The duration of treatment depends on the indication and the patient's response. It is advisable to review the treatment after 2 weeks.                                                                                                                                                                                                                                                                                                                                                                                                         |
| Use in children                      | : No dosage recommendations and indication have been established to date for the use of Ammi Votara in children. So keep out of reach of the children.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| <b>Contraindications</b>             | : Patients who have previously shown a hypersensitivity reaction to diclofenac, aspirin and other NSAIDs, as well as to propylene glycol, isopropyl alcohol.<br>Do not use in the third trimester of Pregnancy.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| <b>Warnings and Precautions</b>      |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Cardiovascular-<br>Thrombotic Events | : Observational studies have indicated that non-selective NSAIDs may be associated with an increased risk of serious cardiovascular events, principally myocardial infarction, which may increase with dose or duration of use. Patients with cardiovascular disease or cardiovascular risk of an adverse cardiovascular event in patient taking NSAIDs, especially in those with cardiovascular risk factors, the lowest effective dose should be used for the shortest possible duration. There is no consistent evidence that the concurrent use of aspirin mitigates the possible increased risk of serious cardiovascular thrombotic events associated with NSAIDs use.                                                                                              |
| Hypertension                         | : NSAIDs may lead to the onset of new hypertension or worsening the pre-existing hypertension and patients taking antihypertensive with NSAIDs may have an impaired antihypertensive response. Caution is advised when prescribing NSAIDs to patients with hypertension. Blood pressure should be monitored closely during initiation of NSAIDs treatment and at regular intervals thereafter.                                                                                                                                                                                                                                                                                                                                                                            |
| Gastrointestinal Events              | : All NSAIDs can cause gastrointestinal discomfort and rarely serious, potentially fatal gastrointestinal effects such as ulcers, bleeding and perforation which may increase with dose or duration of use, but can occur at any time without warning. Caution is advised in patients with risk factors for gastrointestinal events e.g. the elderly, those with a history of serious gastrointestinal events, smoking and alcoholism. When gastrointestinal bleeding or ulcerations occur in patients receiving NSAIDs, The drug should be withdrawn immediately. Doctors should warn patient about signs and symptoms of serious gastrointestinal toxicity. The concurrent use of aspirin and NSAIDs also increase the risk of serious gastrointestinal adverse events. |

|                            |                                                                                                                                                                                                                                                                                                                                         |
|----------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Heart Failure              | : Fluid retention and oedema have been observed in some patients taking NSAIDs, there caution is advised in patients with fluid retention or heart failure.                                                                                                                                                                             |
| Severe cutaneous reactions | : Severe cutaneous reactions, including Stevens - Johnson syndrome and toxic epidermal necrolysis (Lyell's syndrome), have been reported with diclofenac sodium. Patients treated with diclofenac sodium should be closely monitored for signs of hypersensitivity reactions. Discontinue diclofenac sodium immediately if rash occurs. |
| Adverse effects            | : Dermatological : Occasional rashes or skin eruptions. Cases of hair loss, bullous eruptions erythema multiforme, Stevens-Johnson Syndrome, toxic epidermal necrolysis (Lyell's syndrome), and photosensitivity reactions have been reported.                                                                                          |

**Side effects** are ranked under heading of frequency, the most frequent first, using the following convention: very common, common, uncommon, rare, very rare and not known: cannot be estimated from the available data.

|                                                 |                                                                             |
|-------------------------------------------------|-----------------------------------------------------------------------------|
| Immune system disorder;                         |                                                                             |
| Very rare                                       | Hypersensitivity (including urticaria), angioneurotic oedema.               |
| Infections and infestations;                    |                                                                             |
| Very rare                                       | Rash pustular.                                                              |
| Respiratory, thoracic and mediastinal disorders |                                                                             |
| Very rare                                       | Asthma.                                                                     |
| Skin and subcutaneous tissue disorders          |                                                                             |
| Common                                          | Rash, eczema, erythema, dermatitis (including dermatitis contact), pruritus |
| Rare                                            | Dermatitis bullous                                                          |
| Very rare                                       | Photosensitivity reaction                                                   |
| Not known                                       | Burning sensation at the application site, Dry skin                         |

Although less likely with the topical administration, some side effects normally associated with systemically administered diclofenac may also occur. The prolonged use of diclofenac in a relatively extensive area can cause systemic side effects such as nausea, vomiting, diarrhoea or epigastric pain.

Topical preparations of diclofenac gel may cause application site reactions. It should be applied only to intact skin surfaces, and not to broken skin or open wounds. It should not be allowed to come into contact with the eyes or with mucous membranes.

|                                                    |                                                                                                                                                                                              |
|----------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Pregnancy and Lactation</b>                     | : No controlled studies are available in pregnant woman and lactation. Avoid the use of Ammi Votara Emulsion Gel and do not use in the third trimester of Pregnancy.                         |
| <b>Symptoms and Treatment of Overdose</b>          | : Significant systemic reactions resulting from improper use on accidental overdosage (e.g. in children) should be treated with the usual measures employed to manage poisoning with NSAIDs. |
| <b>Interactions with Other Medicament</b>          | : No drug interactions during treatment with Ammi Votara Emulsion Gel have yet been reported.                                                                                                |
| <b>Effects on Ability to Drive and Use Machine</b> | : Cutaneous application of topical diclofenac has no or negligible influence on the ability to drive and use machines.                                                                       |
| <b>Storage conditions</b>                          | : Store at room temperature not exceeding 30°C.                                                                                                                                              |
| <b>Shelf-life</b>                                  | : 3 years                                                                                                                                                                                    |
| <b>Pack size</b>                                   | : Laminated aluminium tube of 20 g.                                                                                                                                                          |

Product Registration Holder :  
**Healoi Pharmaceuticals Sdn. Bhd.,**  
 74-3, Jalan Wangsa Delima 6, KLSC Wangsa Maju, 53300 Kuala Lumpur, Malaysia

Manufactured by :  
**MacroPhar Co., Ltd.**  
 89 Soi Pattanakarn 20 Yaek 4, Pattanakarn Road, Suan Luang, Suan Luang, Bangkok 10250, Thailand.  
 Tel: (662) 314-6671 Fax: (662) 318 6091

Date of revision of Package Insert : 05/09/2024