



GENTACIN CREAM

Description: White colour cream.

Content:

Gentamicin sulphate eq. to gentamicin.....0.10%w/w
Preservative: Chlorocresol 0.1%w/w

Use of topical gentamicin in closed hospital settings is actively discouraged.

Indication:

For the treatment of primary and secondary skin infections caused by susceptible bacteria.

Pharmacology:

Gentamicin sulphate is a broad spectrum aminoglycoside antibiotic which is effective against a wide range of Gram + ve and Gram - ve bacteria, including Staphylococcus aureus (coagulase + ve, coagulase - ve and some penicillinase-producing strains), Streptococci group A (α and β hemolytic), species of Pseudomonas, Proteus vulgaris, E-coli, Aerobacter aerogenes, Klebsiella pneumoniae and many coliform bacilli. It is bactericidal and acts by binding to a specific receptor protein on the 30S subunit of bacterial ribosomes and interfere with an initiation complex between mRNA and the 30S subunit inhibiting protein synthesis. DNA may be misread, thus producing non-functional proteins, polyribosomes are split apart and unable to synthesize proteins.

Systemic absorption of gentamicin has been reported after topical use on denuded skin and burns. Metabolic reactions of gentamicin is by conjugation. Binding of gentamicin to plasma proteins is usually low.

Adverse effects:

Topical application of gentamicin may give rise to allergy e.g. redness, swelling, itching or other signs of irritation not present before therapy.

Precautions:

Excessive or prolonged use of gentamicin may cause overgrowth of lesions by fungi or non-susceptible bacteria. Use of topical antibacterials may lead to skin sensitization, resulting in hypersensitivity reactions with subsequent topical use of the medication.

Pregnancy & Lactation:

Risk benefits should be considered as safety of use has not been established.

Drug Interactions:

Nil for topical application.

Contraindications:

It is contraindicated in patients hypersensitive to gentamicin or any related components.

Dosage and administration:

Adults and children over 1 year of age : Apply to affected skin two times to three times daily.

Symptoms & treatment of overdose:

Subsequent topical use of gentamicin may result in hypersensitivity reactions e.g. itching, redness, swelling or other signs of irritation. If the symptoms above occur, discontinue use and wash the affected part with water. Other symptomatic treatment may also be applied.

Storage Conditions : Keep container tightly closed. Store in a cool, dry place, below 30°C. Protect from light. Keep away from children.

Presentation : 15gm/tube

Shelflife : 3 years.

Product registration holder : **PRIME PHARMACEUTICAL SDN. BHD.**

& Manufacturer : 1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah,
14000 Bukit Mertajam, Penang, Malaysia.

28/09/2025