



T3 Mycin CLINDAMYCIN LOTION

NAME AND STRENGTH OF ACTIVE SUBSTANCE : Clindamycin

Phosphate 1.2% w/v (equiv. to Clindamycin 1.0% w/v).

PRODUCT DESCRIPTION : A clear solution.

Other ingredients : Methyl Paraben, Propyl Paraben, Propylene Glycol, Isopropyl Alcohol, Sodium Hydroxide and Purified Water.

PHARMACODYNAMICS : Clindamycin Phosphate, a lincosamide antibiotic, is a chlorinated derivative of the antibiotic lincomycin, and is used in the treatment of serious anaerobic infections.

PHARMACOKINETICS : Following multiple topical applications of clindamycin phosphate at a concentration equivalent to 10mg clindamycin per ml in an isopropyl alcohol and water solution, very low levels of clindamycin are present in the serum (0-3ng/ml) and less than 0.2% of the dose is recovered in urine as clindamycin. Clindamycin activity has been demonstrated in comedones from acne patients. The mean concentration of antibiotic activity in extracted comedones after application of Clindamycin phosphate topical solution for 4 weeks was 597 mcg/g of comedonal material (range 0-1490). Clindamycin in vitro inhibits all *Propionibacterium* acnes cultures tested (MICs 0.4 mcg/ml). Free fatty acids on the skin surface have been reduced from approximately 14% to 2% following application of clindamycin.

INDICATIONS : For the treatment of acne vulgaris.

CONTRAINDICATIONS : History of hypersensitivity to preparations containing clindamycin, lincomycin or propylene glycol, history of regional enteritis or ulcerative colitis, or a history of antibiotic-associated colitis.

DRUG INTERACTIONS : Clindamycin has been shown to have neuromuscular blocking properties that may enhance the action of other neuromuscular blocking agents. Therefore, it should be used with caution in patients receiving such agents. Cross resistance has been demonstrated between clindamycin and lincomycin. Antagonism has been demonstrated between clindamycin and erythromycin.

ADVERSE EFFECTS : Orally and parenterally administered clindamycin has been associated with severe colitis which may end fatally. Cases of diarrhoea, bloody diarrhoea and colitis (including pseudomembranous colitis) have been reported as adverse reactions in patients treated with topical formulations of clindamycin. Other local effects include contact dermatitis, irritation (e.g erythema, peeling, and burning), oily skin and gram-negative folliculitis.

PRECAUTIONS/WARNINGS : Diarrhoea, bloody diarrhoea and colitis (including pseudomembranous colitis) have been reported with the use of topical clindamycin. When significant diarrhoea

occurs, the drug should be discontinued. Avoid contact with eyes. **Usage in pregnancy :** Safety for use in pregnancy has not been established.

Usage in the Newborn and Infants : Appropriate studies on the relationship of age to the effect of topical clindamycin have not been performed in children up to 12 years of age. Safety and efficacy have not been established.

Nursing Mothers : Clindamycin has been reported to appear in the breast milk in ranges 0.7 to 3.8 mg/ml; therefore it is not recommended for nursing mothers.

- Clindamycin therapy has been associated with severe colitis which may end fatally.
- It should be reserved for serious infections where less toxic antimicrobial agents are inappropriate.
- It should not be used in patients with nonbacterial infections, such as most upper respiratory tract infections.
- Its use in newborns is contraindicated.

ROUTE OF ADMINISTRATION : Topical.

RECOMMENDED DOSAGE : Apply to the affected skin area twice daily.

HOW TO USE : 1. Clean and dry skin area to be treated. 2. Apply a thin film twice daily to the affected area. Avoid eyes and mouth. If medication accidentally enters eyes, rinse thoroughly with water. 3. Use dabbing motion with the tip rather than rolling motion. If tip dries out, invert the bottle and depress tip several times until it becomes moist.

SYMPTOMS AND TREATMENT FOR OVERDOSE AND ANTIDOTE(S) : Pseudomembranous colitis has been reported. Mild cases usually respond to discontinuation of the drug alone. In moderate to severe cases, consideration should be given to management with fluids and electrolytes, protein supplementation and treatment with antibacterial drug clinically effective against *Clostridium difficile*.

PACKING : 30ml frosted glass bottle with a special applicator designed so that the solution may be applied directly to the affected skin area.

STORAGE : Store below 30°C. For external use only. Keep out of reach of children.

Recommended shelf-life : 3 years.

MANUFACTURED BY :

HOE Pharmaceuticals Sdn. Bhd.

Lot 10, Jalan Sultan Mohamed 6, Bandar Sultan Suleiman, 42000 Port Klang, Selangor, MALAYSIA.

Revision date : April 2023